

# On nonacyclicity of the quotient space of $\mathbb{R}^3$ by the solenoid

Umed H. Karimov<sup>a</sup>, Dušan Repovš<sup>b,\*</sup>

<sup>a</sup> *Institute of Mathematics, Academy of Sciences of Tajikistan Ul. Ainy, 299<sup>A</sup>, Dushanbe 734063, Tajikistan*

<sup>b</sup> *Institute of Mathematics, Physics and Mechanics, University of Ljubljana, P.O. Box 2964, Ljubljana 1001, Slovenia*

Received 29 May 2001; received in revised form 2 May 2002

---

## Abstract

It is well-known that the quotient space of the 3-dimensional Euclidean space  $\mathbb{R}^3$  by the dyadic solenoid is not simply connected. We prove that the singular homology of this quotient space is uncountable.

© 2003 Elsevier B.V. All rights reserved.

*MSC:* primary 54B15; secondary 55N10, 55Q52

*Keywords:* Fundamental group; Simple connectivity; Solenoid; Projective telescope; Hawaiian earrings

---

## 1. Introduction

Bing [1] was the first to observe that the quotient space  $\mathbb{R}^3/\Sigma_2$  of the 3-dimensional Euclidean space  $\mathbb{R}^3$  by the dyadic solenoid  $\Sigma_2$  has a nontrivial fundamental group (a complete proof of this result was first published in [8,9]). However, not much is known about its properties. Therefore it is of interest to understand the nature of this group.

The quotient space  $\mathbb{R}^3/\Sigma_2$  is homotopy equivalent to the dyadic projective telescope  $\mathcal{P}_2\mathcal{T}$ . Bogley and Sieradski have shown that the fundamental group  $\pi_1(\mathcal{P}_2\mathcal{T})$  is non-Abelian [2,11]. The purpose of the present paper is to show that the abelianization of the fundamental group  $\pi_1(\mathbb{R}^3/\Sigma_{\mathcal{P}})$  of the quotient space  $\mathbb{R}^3$  by any solenoid  $\Sigma_{\mathcal{P}}$  is an *uncountable* group.

---

\* Corresponding author.

*E-mail addresses:* [umed@ac.tajik.net](mailto:umed@ac.tajik.net) (U.H. Karimov), [dusan.repovs@uni-lj.si](mailto:dusan.repovs@uni-lj.si) (D. Repovš).

**Theorem 1.1.** *The quotient space  $\mathbb{R}^3/\Sigma_{\mathcal{P}}$  of  $\mathbb{R}^3$  by any solenoid  $\Sigma_{\mathcal{P}}$  is homotopy equivalent to the projective telescope  $\mathcal{PT}$  and the singular homology group  $H_1(\mathbb{R}^3/\Sigma_{\mathcal{P}}; \mathbb{Z})$  is uncountable.*

## 2. Preliminaries

Let  $S^1$  be the oriented unit circle in the complex plane  $\mathbb{C}$ . Consider the following inverse sequence  $\mathcal{P}$ :

$$P_0 \xleftarrow{f_0} P_1 \xleftarrow{f_1} P_2 \xleftarrow{f_1} \dots$$

where  $P_0$  is a point,  $P_k$  is the circle  $S^1$  and  $f_k : S^1 \rightarrow S^1$  is the standard continuous mapping of degree  $n_k$ ,  $n_k > 1$ , for every  $k > 0$ . The inverse limit  $\varprojlim \mathcal{P}$  is called the *solenoid*  $\Sigma_{\mathcal{P}}$ . The space  $\Sigma_{\mathcal{P}}$  is one-dimensional, compact and metric. It has a standard embedding into  $\mathbb{R}^3$  (see, e.g., [5, pp. 230–231]). If  $n_k = 2$  for all  $k$ , then  $\Sigma_{\mathcal{P}}$  is called the *dyadic solenoid* and denoted by  $\Sigma_2$ .

Let  $C(f_0, f_1, f_2, \dots)$  be the *infinite mapping cylinder* (see, e.g., [6,7,10]) and let  $\tilde{\mathcal{P}}$  be its natural compactification by the solenoid  $\Sigma_{\mathcal{P}}$ . The projective telescope  $\mathcal{PT}$  is the one-point compactification of  $C(f_0, f_1, f_2, \dots)$  by some point  $\{pt\}$ . We consider  $\{pt\}$  as the base point of  $\mathcal{PT}$  and the circles  $P_k$  for  $k = \{1, 2, 3, \dots\}$  as the natural subspaces of  $\mathcal{PT}$ .

Hereafter, by homology we shall mean the singular homology with integer coefficients. Since the one-dimensional homology group of a path-connected space is the abelianization of the fundamental group, our results strengthen Bing's theorem mentioned above [1,8,9].

To prove Theorem 1.1 we shall need the following results:

**Theorem 2.1** (Borsuk [3,9]). *Let  $W$  be a strong deformation retract of  $\widehat{W}$  and let  $X$  be any continuum in  $W$ . Then  $W/X$  is a strong deformation retract of  $\widehat{W}/X$ . Thus in particular,  $W/X$  and  $\widehat{W}/X$  have the same homotopy type.*

**Proposition 2.2.** *The compactum  $\mathcal{PT}$  is an absolute retract.*

**Proof.** The proposition is a direct consequence of well-known results (see, e.g., [7, p. 104]).  $\square$

Consider the following closed subset of  $S^1$ :

$$A = \left\{ e^{2\pi i t} \in S^1 \mid t = \frac{1}{k}, k \in \mathbb{N} \right\}.$$

The quotient space  $S^1/A$  is homeomorphic to the *Hawaiian earring*  $\mathcal{H}$ , i.e., to the compact bouquet of a countable number of circles  $\{S_k^1\}_{k \in \mathbb{N}}$ .

Let  $p : S^1 \rightarrow \mathcal{H}$  be the canonical projection,  $\mathbb{Z}$  the infinite cyclic group and  $\mathbb{Z}_n$  the finite cyclic subgroup of order  $n$  of  $S^1$ :

$$\mathbb{Z}_n = \left\{ e^{2\pi i t} \in S^1 \mid t = \frac{k}{n}, k = 1, 2, \dots, n \right\}.$$

### 3. Proof of Theorem 1.1

Since the space  $\tilde{\mathcal{P}}$  is a 2-dimensional compactum, it can be considered as a closed subspace of  $\mathbb{R}^5$ . Since  $\mathbb{R}^5$  and (by Proposition 2.2)  $\tilde{\mathcal{P}}$  is an absolute retract,  $\tilde{\mathcal{P}}$  is a strong deformation retract of  $\mathbb{R}^5$ . The compactum  $\Sigma_{\mathcal{P}}$  is a subset of  $\tilde{\mathcal{P}}$ , therefore by Theorem 2.1 the quotient space  $\mathbb{R}^5/\Sigma_{\mathcal{P}}$  is homotopy equivalent to the quotient space  $\tilde{\mathcal{P}}/\Sigma_{\mathcal{P}}$ , which is obviously homeomorphic to the projective telescope  $\mathcal{PT}$ .

Since the homotopy type of  $\mathbb{R}^5/\Sigma_{\mathcal{P}}$  does not depend on the way in which  $\Sigma_{\mathcal{P}}$  is embedded into  $\mathbb{R}^5$  (see Theorem 1 in [9]), we can assume that  $\Sigma_{\mathcal{P}}$  is embedded into  $\mathbb{R}^5$  as the composition of the standard embeddings  $\Sigma_{\mathcal{P}} \subset \mathbb{R}^3 \times \{0\} \subset \mathbb{R}^3 \times \mathbb{R}^2$ , where 0 is the origin of  $\mathbb{R}^2$ . By Theorem 2.1,  $\mathbb{R}^3/\Sigma_{\mathcal{P}}$  is homotopy equivalent to  $\mathbb{R}^5/\Sigma_{\mathcal{P}}$  and therefore to the projective telescope  $\mathcal{PT}$ . The first part of Theorem 1.1 is thus proved.

Suppose now that to the contrary,  $H_1(\mathcal{PT})$  were a countable group. Consider  $\mathcal{PT}$  as the union:  $\mathcal{PT} = C(f_0) \cup C(f_1, f_2, f_3, \dots)^*$ , where  $C(f_0)$  is the cylinder of the constant mapping  $f_0: S^1 \rightarrow S^1$  and therefore is a contractible space, and  $C(f_1, f_2, f_3, \dots)^*$  is the one-point compactification of the infinite mapping cylinder  $C(f_1, f_2, f_3, \dots)$ . The intersection of these two subspaces of  $\mathcal{PT}$  is the circle  $S^1$ . Thus it follows by the Mayer–Vietoris exact sequence:

$$\rightarrow H_1(S^1) \rightarrow H_1(C(f_0)) \oplus H_1(C(f_1, f_2, f_3, \dots)^*) \rightarrow H_1(\mathcal{PT}) \rightarrow \dots$$

that the group

$$H_1(C(f_1, f_2, f_3, \dots)^*) \text{ is countable.} \tag{3.1}$$

Consider now  $C(f_{n+1}, f_{n+2}, f_{n+3}, \dots)^*$  as a subspace of  $C(f_1, f_2, f_3, \dots)^*$ . Let  $X_n$  and  $p_n: C(f_1, f_2, f_3, \dots)^* \rightarrow X_n$  be the corresponding quotient space and the quotient mapping. For every sequence of units and zeros  $\alpha = (\alpha_1, \alpha_2, \alpha_3, \dots)$ , let  $g_\alpha: \mathcal{H} \rightarrow \mathcal{H}$  be the mapping such that

$$g_\alpha|_{S^1_k} = \begin{cases} \text{the identity mapping onto its image,} & \text{if } \alpha_k = 1, \\ \text{the constant mapping into the base point,} & \text{if } \alpha_k = 0. \end{cases}$$

Let  $g$  be a mapping of  $\mathcal{H}$  to  $C(f_1, f_2, f_3, \dots)^*$  which maps the base point of  $\mathcal{H}$  to the base point  $\{pt\}$  of  $C(f_1, f_2, f_3, \dots)^*$  and such that the restriction  $g|_{S^1_k}$  only wraps once around the circle  $P_k$  in the positive direction.

The set  $\{g_\alpha\}$  is uncountable. However, the group  $H_1(C(f_1, f_2, f_3, \dots)^*)$  is countable (3.1). Therefore there exist two sequences  $\alpha$  and  $\beta$  such that  $\alpha \neq \beta$  and such that for the mappings  $S^1 \xrightarrow{p} \mathcal{H} \xrightarrow{g_\alpha} \mathcal{H} \xrightarrow{g} C(f_1, f_2, f_3, \dots)^*$  and  $S^1 \xrightarrow{p} \mathcal{H} \xrightarrow{g_\beta} \mathcal{H} \xrightarrow{g} C(f_1, f_2, f_3, \dots)^*$  we obtain the same homomorphism of the corresponding homology groups:

$$(gg_\alpha p)_1 = (gg_\beta p)_1: H_1(S^1) \rightarrow H_1(C(f_1, f_2, f_3, \dots)^*). \tag{3.2}$$

On the other hand, let  $m$  be the minimal number such that  $\alpha_m \neq \beta_m$ . To the projection  $p_m: C(f_1, f_2, f_3, \dots)^* \rightarrow X_m$  there correspond two homomorphisms of homology groups:  $H_1(S^1) \xrightarrow{(p_m g g_\alpha p)_1} H_1(X_m)$  and  $H_1(S^1) \xrightarrow{(p_m g g_\beta p)_1} H_1(X_m)$ . Since  $\alpha_k = \beta_k$  for  $k < m$  and  $\alpha_m \neq \beta_m$ , by construction we have  $(p_m g g_\alpha p)_1(1) \neq (p_m g g_\beta p)_1(1)$ , contradicting (3.2).

**Question 3.1.** Let  $X$  be the Case–Chamberlin continuum [4]. Is then the homology of quotient space  $H_1(\mathbb{R}^3/X)$  nontrivial?

### Acknowledgements

We acknowledge the support by the Ministry for Education, Science and Sport of the Republic of Slovenia research program No. 0101-509 and research grants No. SLO-KIT-04-14-2002 and No. SLO-US-2002-01. The first author thanks Professor B.U. Makhmadaliev for the support during the work on this paper. We thank the referee for comments and suggestions.

### References

- [1] R.H. Bing, Conditions under which monotone decomposition of  $E^3$  are simple connected, *Bull. Amer. Math. Soc.* 63 (1957) 143, Abstract No. 325.
- [2] W.A. Bogley, A.J. Sieradski, Omega-groups II: Weighted presentation for omega-groups, Preprint, Univ. of Oregon, Eugene, OR, 1997.
- [3] K. Borsuk, On the homotopy type of some decomposition spaces, *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astron. Phys.* 18 (1970) 235–239.
- [4] J.H. Case, R.E. Chamberlin, Characterization of tree-like continua, *Pacific J. Math.* 10 (1960) 73–84.
- [5] S. Eilenberg, N. Steenrod, *Foundation of Algebraic Topology*, Princeton University Press, Princeton, NJ, 1952.
- [6] U.H. Karimov, D. Repovš, On suspensions of noncontractible compactum of trivial shape, *Proc. Amer. Math. Soc.* 127 (1999) 627–632.
- [7] J. Krasinkiewicz, On a methods of constructing ANR-sets. An application of inverse limits, *Fund. Math.* 92 (1976) 95–112.
- [8] D.R. McMillan, N. Shrikhande, On the simple connectivity of a quotient space, *Glasnik Mat.* 66 (1983) 113–124.
- [9] N. Shrikhande, Homotopy properties of decomposition spaces, *Fund. Math.* 66 (1983) 119–124.
- [10] L.C. Siebenmann, Chapman’s classification of shapes. A proof using collapsing, *Manuscripta Math.* 16 (1975) 373–384.
- [11] A.J. Sieradski, Omega-groups: Group theory for wild topology, Preprint, Univ. of Oregon, Eugene, OR, 1998.