

ORIGINAL PAPER

Parametric nonlinear resonant Robin problems

Nikolaos S. Papageorgiou^{1,2} | Vicențiu D. Rădulescu^{2,3,4}  | Dušan D. Repovš^{2,5} ¹National Technical University, Department of Mathematics, Zografou Campus, Athens 15780, Greece²Institute of Mathematics, Physics and Mechanics, 1000 Ljubljana, Slovenia³Faculty of Applied Mathematics, AGH University of Science and Technology, 30-059 Kraków, Poland⁴Institute of Mathematics“Simion Stoilow” of the Romanian Academy, P.O. Box 1-764, 014700 Bucharest, Romania⁵Faculty of Education and Faculty of Mathematics and Physics, University of Ljubljana, 1000 Ljubljana, Slovenia**Correspondence**

Vicențiu D. Rădulescu, Institute of Mathematics, Physics and Mechanics, 1000 Ljubljana, Slovenia.

Email: vicentiu.radulescu@imfm.si

Funding information

Slovenian Research Agency, Grant/Award Numbers: P1-0292, J1-8131, J1-7025, N1-0064, N1-0083

Abstract

We consider a nonlinear Robin problem driven by the p -Laplacian. In the reaction we have the competing effects of two nonlinearities. One term is parametric, strictly $(p - 1)$ -sublinear and the other one is $(p - 1)$ -linear and resonant at any nonprincipal variational eigenvalue. Using variational tools from the critical theory (critical groups), we show that for all big values of the parameter λ the problem has at least five nontrivial smooth solutions.

KEYWORDS

critical groups, multiple solutions, nonlinear regularity, resonance, Robin boundary condition, strong comparison, truncation

MSC (2010)

35J20, 35J60, 58E05

1 | INTRODUCTION

Let $\Omega \subseteq \mathbf{R}^N$ be a bounded domain with a C^2 -boundary $\partial\Omega$. In this paper, we study the following parametric nonlinear Robin problem:

$$\left\{ \begin{array}{l} -\Delta_p u(z) = \lambda g(z, u(z)) + f(z, u(z)) \text{ in } \Omega, \\ \frac{\partial u}{\partial n_p} + \beta(z)|u|^{p-2}u = 0 \text{ on } \partial\Omega, \end{array} \right. \quad (P_\lambda)$$

where λ is a positive parameter.

In this problem, Δ_p denotes the p -Laplace differential operator defined by

$$\Delta_p u = \operatorname{div}(|Du|^{p-2} Du) \quad \text{for all } u \in W^{1,p}(\Omega), \quad 1 < p < \infty.$$

In the reaction (the right-hand side) of problem (P_λ) , $g(z, x)$ and $f(z, x)$ are Carathéodory functions (that is, for all $x \in \mathbf{R}$, the mappings $z \mapsto g(z, x)$ and $z \mapsto f(z, x)$ are measurable, while for almost all $z \in \Omega$, the mappings $x \mapsto g(z, x)$ and $x \mapsto f(z, x)$ are continuous functions). These two nonlinearities exhibit different growth near $\pm\infty$ and 0. More precisely, for almost all $z \in \Omega$,

$g(z, \cdot)$ is $(p - 1)$ -sublinear both near 0 and near $\pm\infty$, while $f(z, \cdot)$ is $(p - 1)$ -linear near 0 and $\pm\infty$. In fact, we permit resonance at $\pm\infty$ with respect to any nonprincipal variational eigenvalue of $-\Delta_p$ with Robin boundary condition.

The coefficient $\beta(\cdot)$ that appears in the boundary condition is strictly positive. This is needed in order to be able to use strong comparison techniques, which in the case of the p -Laplace differential operator are difficult to apply.

We denote by $\frac{\partial u}{\partial n_p}$ the conormal derivative of u , which is defined by extension of the map

$$C^1(\overline{\Omega}) \ni u \mapsto \frac{\partial u}{\partial n_p} = |Du|^{p-2}(Du, n)_{\mathbb{R}^N} = |Du|^{p-2} \frac{\partial u}{\partial n},$$

with $n(\cdot)$ being the outward unit normal on $\partial\Omega$.

Using variational tools from the critical point theory, together with suitable truncation and strong comparison techniques and Morse theory (critical groups), we show that for big values of the positive parameter λ , problem (P_λ) admits at least five nontrivial smooth solutions.

Multiplicity results proving three solutions theorems for nonresonant Dirichlet p -Laplacian equations were established by Gasinski & Papageorgiou [9], Guo & Liu [13], and Jiu & Su [15], Liu [18]. Resonant p -Laplacian equations were investigated by Gasinski & Papageorgiou [7,8], Mugnai & Papageorgiou [20], Papageorgiou & Rădulescu [22] (Neumann problems), and Papageorgiou & Rădulescu [21] (Robin problems). In all the above works, the resonance was with respect to the principal eigenvalue. Resonance with respect to higher variational eigenvalues was allowed in the recent works of Papageorgiou, Rădulescu & Repovš [25,26], which dealt with nonparametric equations. None of the aforementioned works produces more than three solutions. Abstract methods closely related with the content of this paper have been developed in the recent monograph of Papageorgiou, Rădulescu & Repovš [27].

2 | MATHEMATICAL BACKGROUND AND HYPOTHESES

Let X be a Banach space and let X^* be its topological dual. We denote by $\langle \cdot, \cdot \rangle$ the duality brackets for the pair (X^*, X) . Given $\varphi \in C^1(X, \mathbb{R})$, we say that $\varphi(\cdot)$ satisfies the ‘‘Cerami condition’’ (the ‘‘C-condition’’ for short), if the following property holds:

‘‘Every sequence $\{u_n\}_{n \geq 1} \subseteq X$ such that $\{\varphi(u_n)\}_{n \geq 1} \subseteq \mathbb{R}$ is bounded and $(1 + \|u_n\|_X)\varphi'(u_n) \rightarrow 0$ in X^* as $n \rightarrow \infty$, admits a strongly convergent subsequence’’.

This compactness-type condition on the functional $\varphi(\cdot)$ leads to a deformation theorem, which is the main analytical tool in deriving the minimax theory of the critical values of φ . One of the main results in that theory is the so-called ‘‘mountain pass theorem’’, which we recall here.

Theorem 2.1. *If $\varphi \in C^1(X, \mathbb{R})$ satisfies the PS-condition, $u_0, u_1 \in X$, $\|u_1 - u_0\| > \rho > 0$,*

$$\max\{\varphi(u_0), \varphi(u_1)\} < \inf\{\varphi(u) : \|u - u_0\| = \rho\} = m_\rho$$

and

$$c = \inf_{\gamma \in \Gamma} \max_{0 \leq t \leq 1} \varphi(\gamma(t)) \quad \text{with} \quad \Gamma = \{\gamma \in C([0, 1], X) : \gamma(0) = u_0, \gamma(1) = u_1\},$$

then $c \geq m_\rho$ and c is a critical value of φ (that is, we can find $\hat{u} \in X$ such that $\varphi'(\hat{u}) = 0$ and $\varphi(\hat{u}) = c$).

The following spaces will play a central role in the analysis of problem (P_λ) :

$$W^{1,p}(\Omega), C^1(\overline{\Omega}) \text{ and } L^p(\partial\Omega).$$

We denote by $\|\cdot\|$ the norm of the Sobolev space $W^{1,p}(\Omega)$. We know that

$$\|u\| = \left(\|u\|_p^p + \|Du\|_p^p \right)^{\frac{1}{p}} \quad \text{for all } u \in W^{1,p}(\Omega).$$

The space $C^1(\overline{\Omega})$ is an ordered Banach space with positive (order) cone

$$C_+ = \{u \in C^1(\overline{\Omega}) : u(z) \geq 0 \text{ for all } z \in \overline{\Omega}\}.$$

This cone has a nonempty interior which is given by

$$D_+ = \{u \in C_+ : u(z) > 0 \text{ for all } z \in \overline{\Omega}\}.$$

In fact, D_+ is the interior of C_+ when the latter is furnished with the weaker $C(\overline{\Omega})$ -norm topology.

On $\partial\Omega$ we consider the $(N-1)$ -dimensional Hausdorff (surface) measure $\sigma(\cdot)$. Using this measure on $\partial\Omega$, we can define in the usual way the boundary Lebesgue spaces $L^q(\partial\Omega)$, $1 \leq q \leq \infty$. We know that there exists a unique continuous linear map $\gamma_0 : W^{1,p}(\Omega) \rightarrow L^p(\partial\Omega)$, known as the “trace map”, such that

$$\gamma_0(u) = u|_{\partial\Omega} \text{ for all } u \in W^{1,p}(\Omega) \cap C(\overline{\Omega}).$$

The trace map defines boundary values for all Sobolev functions. We know that $\gamma_0(\cdot)$ is a compact map into $L^q(\partial\Omega)$ for all $q \in \left[1, \frac{(N-1)p}{N-p}\right)$ when $p < N$, and into $L^q(\partial\Omega)$ for all $1 \leq q < \infty$ when $N \leq p$. We have

$$\text{im } \gamma_0 = W^{\frac{1}{p'}, p}(\partial\Omega) \text{ and } \ker \gamma_0 = W_0^{1,p}(\Omega).$$

Recall that p' denotes the conjugate exponent of p (that is, $\frac{1}{p} + \frac{1}{p'} = 1$). In what follows, for the sake of notational simplicity, we drop the use of trace map γ_0 . All restrictions of Sobolev functions on $\partial\Omega$ are understood in the sense of traces.

Our hypotheses on the boundary coefficient $\beta(\cdot)$ are the following:

$$H(\beta): \beta \in C^{0,\alpha}(\partial\Omega) \text{ with } 0 < \alpha < 1 \text{ and } \beta(z) > 0 \text{ for all } z \in \partial\Omega.$$

In the sequel, we denote by $\tau : W^{1,p}(\Omega) \rightarrow \mathbf{R}$ the C^1 -functional defined by

$$\tau(u) = \|Du\|_p^p + \int_{\partial\Omega} \beta(z)|u|^p d\sigma.$$

By Proposition 2.4 of Gasinski & Papageorgiou [11], we know that $\tau(\cdot)^{\frac{1}{p}}$ is an equivalent norm on $W^{1,p}(\Omega)$. So, there exist $c_1, c_2 > 0$ such that

$$c_1 \|u\|^p \leq \tau(u) \leq c_2 \|u\|^p \text{ for all } u \in W^{1,p}(\Omega). \quad (2.1)$$

Let $f_0 : \Omega \times \mathbf{R} \rightarrow \mathbf{R}$ be a Carathéodory function such that

$$|f_0(z, x)| \leq a_0(z)(1 + |x|^{r-1}) \text{ for almost all } z \in \Omega \text{ and all } x \in \mathbf{R},$$

with $a_0 \in L^\infty(\Omega)$, $1 < r \leq p^*$, where p^* is the Sobolev critical exponent corresponding to p , hence

$$p^* = \begin{cases} \frac{Np}{N-p} & \text{if } p < N, \\ +\infty & \text{if } N \leq p. \end{cases}$$

We set $F_0(z, x) = \int_0^x f_0(z, s) ds$ and consider the C^1 -functional $\varphi_0 : W^{1,p}(\Omega) \rightarrow \mathbf{R}$ defined by

$$\varphi_0(u) = \frac{1}{p} \tau(u) - \int_{\Omega} F_0(z, u) dz \text{ for all } u \in W^{1,p}(\Omega).$$

The next proposition is a special case of a more general result of Papageorgiou & Rădulescu [23]. The proposition is essentially an outgrowth of the nonlinear regularity theory of Lieberman [17].

Proposition 2.2. *Assume that $u_0 \in W^{1,p}(\Omega)$ is a local $C^1(\overline{\Omega})$ -minimizer of φ_0 , that is, there exists $\rho_0 > 0$ such that*

$$\varphi_0(u_0) \leq \varphi_0(u_0 + h) \text{ for all } h \in C^1(\overline{\Omega}) \text{ with } \|h\|_{C^1(\overline{\Omega})} \leq \rho_0,$$

Then $u_0 \in C^{1,\alpha}(\overline{\Omega})$ for some $\alpha \in (0, 1)$ and u_0 is also a local $W^{1,p}(\Omega)$ -minimizer of φ_0 , that is, there exists $\rho_1 > 0$ such that

$$\varphi_0(u_0) \leq \varphi_0(u_0 + h) \quad \text{for all } h \in W^{1,p}(\Omega) \text{ with } \|h\| \leq \rho_1.$$

It is well-known that in the nonlinear case ($p \neq 2$), it is difficult to produce strong comparison results and more restrictive conditions are needed on the data of the problem. The next proposition is a special case of a more general result of Gasinski & Papageorgiou [11, Proposition 3.4].

Proposition 2.3. *If $h_1, h_2 \in L^\infty(\Omega)$, $h_1(z) \leq h_2(z)$ for almost all $z \in \Omega$, $h_1 \not\equiv h_2$ and $u_1, u_2 \in C^1(\overline{\Omega})$ satisfy $u_1 \leq u_2$ and*

$$\begin{aligned} -\Delta_p u_1(z) &= h_1(z) \text{ for almost all } z \in \Omega, \quad \left. \frac{\partial u_1}{\partial n} \right|_{\partial\Omega} < 0, \\ -\Delta_p u_2(z) &= h_2(z) \text{ for almost all } z \in \Omega, \quad \left. \frac{\partial u_2}{\partial n} \right|_{\partial\Omega} < 0, \end{aligned}$$

then $u_2 - u_1 \in \text{int } \hat{C}_+ = \left\{ u \in C^1(\overline{\Omega}) : u \Big|_{\Omega} > 0, \left. \frac{\partial u}{\partial n} \right|_{u^{-1}(0)} < 0 \right\}$.

Let $A : W^{1,p}(\Omega) \rightarrow W^{1,p}(\Omega)^*$ be the nonlinear map defined by

$$\langle A(u), h \rangle = \int_{\Omega} |Du|^{p-2} (Du, Dh)_{\mathbf{R}^N} dz \quad \text{for all } u, h \in W^{1,p}(\Omega).$$

The next proposition is a special case of Problem 2.192 of Gasinski & Papageorgiou [10, p. 279].

Proposition 2.4. *The map $A(\cdot)$ is bounded (that is, it maps bounded sets to bounded sets), continuous, monotone (thus, maximal monotone, too) and of type $(S)_+$, that is,*

$$\text{if } u_n \xrightarrow{w} u \text{ in } W^{1,p}(\Omega) \text{ and } \limsup_{n \rightarrow \infty} \langle A(u_n), u_n - u \rangle \leq 0, \text{ then } u_n \rightarrow u \text{ in } W^{1,p}(\Omega).$$

We will need some basic facts about the spectrum of the negative p -Laplacian with Robin boundary condition. So, we consider the following nonlinear eigenvalue problem:

$$\left\{ \begin{array}{l} -\Delta_p u(z) = \hat{\lambda} |u(z)|^{p-2} u(z) \quad \text{in } \Omega, \\ \left. \frac{\partial u}{\partial n_p} + \beta(z) |u|^{p-2} u = 0 \right. \quad \text{on } \partial\Omega. \end{array} \right\} \tag{2.2}$$

We say that $\hat{\lambda} \in \mathbf{R}$ is an ‘‘eigenvalue’’ of (2.2), if the problem admits a nontrivial solution $\hat{u} \in W^{1,p}(\Omega)$, known as an eigenfunction corresponding to $\hat{\lambda}$. The nonlinear regularity theory of Lieberman [17, Theorem 2], implies that $\hat{u} \in C^1(\overline{\Omega})$. There is a smallest eigenvalue $\hat{\lambda}_1$ which has the following properties:

- $\hat{\lambda}_1$ is isolated (that is, we can find $\varepsilon > 0$ such that the open interval $(\hat{\lambda}_1, \hat{\lambda}_1 + \varepsilon)$ contains no eigenvalues);
- $\hat{\lambda}_1$ is simple (that is, if \hat{u}, \tilde{u} are eigenfunctions corresponding to $\hat{\lambda}_1$, then $\hat{u} = \xi \tilde{u}$ for some $\xi \in \mathbf{R} \setminus \{0\}$);
- we have

$$\hat{\lambda}_1 = \inf \left\{ \frac{\tau(u)}{\|u\|_p^p} : u \in W^{1,p}(\Omega), u \neq 0 \right\} > 0 \quad (\text{see (2.1)}). \tag{2.3}$$

The infimum in (2.3) is realized on the corresponding one-dimensional eigenspace. From the above properties it follows that the elements of this eigenspace do not change sign and they, of course, belong in $C^1(\overline{\Omega})$. Let \hat{u}_1 denote the positive, L^p -normalized (that is, $\|\hat{u}_1\|_p = 1$) eigenfunction corresponding to $\hat{\lambda}_1$. We have $\hat{u}_1 \in C_+ \setminus \{0\}$ and in fact, by the nonlinear Hopf’s boundary point theorem (see Gasinski & Papageorgiou [6, p. 738]), we have $\hat{u}_1 \in D_+$.

Let $\hat{\sigma}(p)$ denote the set of eigenvalues of (2.2). It is easy to check that the set $\hat{\sigma}(p) \subseteq (0, +\infty)$ is closed. So, the second eigenvalue of (2.2) is well-defined by

$$\hat{\lambda}_2 = \min \{ \hat{\lambda} \in \hat{\sigma}(p) : \hat{\lambda} \neq \hat{\lambda}_1 \}.$$

The Ljusternik–Schnirelmann minimax scheme gives us in addition to $\hat{\lambda}_1$ and $\hat{\lambda}_2$, a whole strictly increasing sequence $\{\hat{\lambda}_k\}_{k \in \mathbf{N}}$ of distinct eigenvalues of (2.2) such that $\hat{\lambda}_k \rightarrow +\infty$. These are known as “variational eigenvalues”. Depending on the index used in the Ljusternik–Schnirelmann minimax scheme, we produce a corresponding sequence of variational eigenvalues. We know that these sequences coincide in the first two elements. However, we do not know if the variational eigenvalues are independent of the index used or they exhaust $\hat{\sigma}(p)$. This is the case if $p = 2$ (linear eigenvalues problem). Here we consider the sequence of variational eigenvalues generated by the Fadell–Rabinowitz cohomological index (see [5]). In this way we can use the results of Cingolani & Degiovanni [4] (see also Papageorgiou, Rădulescu & Repovš [25, Proposition 12]). Note that if $\hat{\lambda} \neq \hat{\lambda}_1$, then the eigenfunctions are sign-changing.

The following lemma is a simple consequence of the above properties of $\hat{\lambda}_1 > 0$ (see Papageorgiou, Rădulescu & Repovš [25, Lemma 14]).

Lemma 2.5. *If $\vartheta \in L^\infty(\Omega)$, $\vartheta(z) \leq \hat{\lambda}_1$ for almost all $z \in \Omega$, $\vartheta \not\equiv \hat{\lambda}_1$, then there exists $c_3 > 0$ such that*

$$c_3 \|u\|^p \leq \tau(u) - \int_{\Omega} \vartheta(z)|u|^p dz \quad \text{for all } u \in W^{1,p}(\Omega).$$

Next, we recall some basic definitions and facts from the theory of critical groups. So, let X be a Banach space and let $\varphi \in C^1(X, \mathbf{R})$, $c \in \mathbf{R}$. We introduce the following sets:

$$K_\varphi = \{u \in X : \varphi'(u) = 0\}, \quad K_\varphi^c = \{u \in K_\varphi : \varphi(u) = c\}, \quad \varphi^c = \{u \in X : \varphi(u) \leq c\}.$$

Given a topological pair (Y_1, Y_2) such that $Y_2 \subseteq Y_1 \subseteq X$, we denote by $H_k(Y_1, Y_2)$ ($k \in \mathbf{N}_0$) the k th relative singular homology group with integer coefficients. Recall that $H_k(Y_1, Y_2) = 0$ for all $k \in -\mathbf{N}$. Suppose that $u \in K_\varphi^c$ is isolated. The critical groups of φ at u are defined by

$$C_k(\varphi, u) = H_k(\varphi^c \cap U, \varphi^c \cap U \setminus \{u\}) \quad \text{for all } k \in \mathbf{N}_0,$$

with U a neighborhood of u such that $K_\varphi \cap \varphi^c \cap U = \{u\}$. The excision property of singular homology implies that this definition is independent of the choice of the isolating neighborhood U .

Suppose that φ satisfies the C-condition and $\inf \varphi(K_\varphi) > -\infty$. Then the critical groups of φ at infinity are defined by

$$C_k(\varphi, \infty) = H_k(X, \varphi^c) \quad \text{for all } k \in \mathbf{N}_0,$$

with $c < \inf \varphi(K_\varphi)$. This definition is independent of the choice of the level $c < \inf \varphi(K_\varphi)$. Indeed, suppose that $c' < c < \inf \varphi(K_\varphi)$. Then the second deformation theorem (see, for example, Gasinski & Papageorgiou [6, p. 628]) implies that $\varphi^{c'}$ is a strong deformation retract of φ^c . Therefore

$$H_k(X, \varphi^c) = H_k(X, \varphi^{c'}) \quad \text{for all } k \in \mathbf{N}_0$$

(see Motreanu, Motreanu & Papageorgiou [19, Corollary 6.15, p. 145]).

Assume that $\varphi \in C^1(X, \mathbf{R})$ satisfies the C-condition and that K_φ is finite. We introduce the following items:

$$M(t, u) = \sum_{k \geq 0} \text{rank } C_k(\varphi, u) t^k \quad \text{for all } t \in \mathbf{R}, u \in K_\varphi,$$

$$P(t, \infty) = \sum_{k \geq 0} \text{rank } C_k(\varphi, \infty) t^k \quad \text{for all } t \in \mathbf{R}.$$

Then the Morse relation says that there exists $Q(t) = \sum_{k \geq 0} \hat{\beta}_k t^k$ a formal series in $t \in \mathbf{R}$ with nonnegative integer coefficients $\hat{\beta}_k$ such that

$$\sum_{u \in K_\varphi} M(t, u) = P(t, \infty) + (1+t)Q(t) \quad \text{for all } t \in \mathbf{R}. \quad (2.4)$$

Now let us fix some basic notation which we will use throughout this work. So, for $x \in \mathbf{R}$, we set $x^\pm = \max\{\pm x, 0\}$. Then for $u \in W^{1,p}(\Omega)$ we define

$$u^\pm(\cdot) = u(\cdot)^\pm.$$

We know that

$$u^\pm \in W^{1,p}(\Omega), \quad u = u^+ - u^-, \quad |u| = u^+ + u^-.$$

If $u, v \in W^{1,p}(\Omega)$ and $v \leq u$, then by $[v, u]$ we denote the ordered interval in $W^{1,p}(\Omega)$ defined by

$$[v, u] = \{y \in W^{1,p}(\Omega) : v(z) \leq y(z) \leq u(z) \text{ for almost all } z \in \Omega\}.$$

By $\text{int}_{C^1(\bar{\Omega})}[v, u]$, we denote the interior in the $C^1(\bar{\Omega})$ -norm topology of $[v, u] \cap C^1(\bar{\Omega})$. We also define

$$[u) = \{y \in W^{1,p}(\Omega) : u(z) \leq y(z) \text{ for almost all } z \in \Omega\}.$$

For $u, v \in W^{1,p}(\Omega)$ with $v(z) \neq 0$ for almost all $z \in \Omega$, we define

$$R(u, v)(z) = |Du(z)|^p - |Dv(z)|^{p-2} \left(Dv(z), D\left(\frac{u^p}{v^{p-1}}\right)(z) \right)_{\mathbf{R}^N}, \quad z \in \Omega.$$

From the nonlinear Picone's identity of Allegretto & Huang [2], we have the following property.

Proposition 2.6. *If $u, v : \Omega \rightarrow \mathbf{R}$ are differentiable functions with $u(z) \geq 0$ and $v(z) > 0$ for all $z \in \Omega$, then $R(u, v)(z) \geq 0$ for almost all $z \in \Omega$ and equality holds if and only if $u = \xi v$ with $\xi \geq 0$.*

Finally, if $k, m \in \mathbf{N}_0$, then by $\delta_{k,m}$ we denote the Kronecker symbol, that is,

$$\delta_{k,m} = \begin{cases} 1 & \text{if } k = m, \\ 0 & \text{if } k \neq m. \end{cases}$$

Next, we introduce our hypotheses on the two nonlinearities in the reaction of (P_λ) .

$H(g) : g : \Omega \times \mathbf{R} \rightarrow \mathbf{R}$ is a Carathéodory function such that $g(z, 0) = 0$ for almost all $z \in \Omega$ and

(i) for every $\rho > 0$, there exists $a_\rho \in L^\infty(\Omega)$ such that

$$|g(z, x)| \leq a_\rho(z) \text{ for almost all } z \in \Omega \text{ and all } |x| \leq \rho,$$

$$0 < g(z, x)x \text{ for almost all } z \in \Omega \text{ and all } x \in \mathbf{R} \setminus \{0\};$$

(ii) $\lim_{x \rightarrow 0} \frac{g(z,x)}{|x|^{p-2}x} = 0$ and there exists $1 < \tau < p$ such that $\lim_{x \rightarrow \pm\infty} \frac{g(z,x)}{|x|^{\tau-2}x} = 0$ uniformly for almost all $z \in \Omega$.

$H(f) : f : \Omega \times \mathbf{R} \rightarrow \mathbf{R}$ is a Carathéodory function such that $f(z, 0) = 0$ for almost all $z \in \Omega$ and

(i) for every $\rho > 0$, there exists $\hat{a}_\rho \in L^\infty(\Omega)$ such that

$$|f(z, x)| \leq \hat{a}_\rho(z) \text{ for almost all } z \in \Omega \text{ and all } |x| \leq \rho;$$

(ii) $\lim_{x \rightarrow \pm\infty} \frac{f(z,x)}{|x|^{p-2}x} = \hat{\lambda}_m$ uniformly for almost all $z \in \Omega$ for some $m \in \mathbf{N}, m \geq 2$ and if $F(z, x) = \int_0^x f(z, s) ds$ then $\liminf_{x \rightarrow \pm\infty} \frac{\rho F(z,x) - f(z,x)x}{|x|^\tau} > 0$ uniformly for almost all $z \in \Omega$;

(iii) there exists $\vartheta \in L^\infty(\Omega)$ such that

$$\vartheta(z) \leq \hat{\lambda}_1 \text{ for almost all } z \in \Omega, \vartheta \neq \hat{\lambda}_1,$$

$$\limsup_{x \rightarrow 0} \frac{f(z, x)}{|x|^{p-2}x} \leq \vartheta(z) \text{ uniformly for almost all } z \in \Omega.$$

$H_0 : \text{For almost all } z \in \Omega \text{ and every } \lambda > 0, \text{ the mapping } x \mapsto \lambda g(z, x) + f(z, x) \text{ is strictly increasing.}$

Remark 2.7. Hypothesis $H(g)(ii)$ implies that $g(z, \cdot)$ is strictly sublinear near $\pm\infty$ and 0. On the other hand, hypothesis $H(f)(ii)$ implies that $f(z, \cdot)$ is $(p - 1)$ -linear near $\pm\infty$. Note that hypotheses $H(g)(ii), H(f)(ii)$ imply that problem (P_λ) at $\pm\infty$ is

resonant with respect to a nonprincipal variational eigenvalue of the Robin p -Laplacian. Clearly, the above hypotheses imply that

$$|g(z, x)|, |f(z, x)| \leq c_4(1 + |x|^{p-1}) \text{ for almost all } z \in \Omega \text{ and all } x \in \mathbf{R} \quad (2.5)$$

with $c_4 > 0$. In the sequel, we shall denote $G(z, x) = \int_0^x g(z, s) ds$.

Example 2.8. The following functions satisfy hypotheses $H(g)$, $H(f)$. For the sake of simplicity, we drop the z -dependence.

$$g(x) = \begin{cases} |x|^{r-2}x & \text{if } |x| \leq 1, \\ |x|^{s-2}x & \text{if } 1 < |x|, \end{cases} \quad 1 < s < p < r;$$

$$f(x) = \begin{cases} \vartheta|x|^{p-2}x & \text{if } |x| \leq 1, \\ \hat{\lambda}_m|x|^{p-2}x + (\hat{\lambda}_m - \vartheta)|x|^{q-2}x & \text{if } 1 < |x|, \end{cases} \quad s < q < p, \vartheta < \hat{\lambda}_1.$$

3 | SOLUTIONS OF CONSTANT SIGN

On account of hypotheses $H(g)(ii)$, $H(f)(ii)$ and (2.5), we see that given $\lambda > 0$, $\varepsilon > 0$ and $r \in (p, p^*)$, we can find $c_5 > 0$ such that

$$[\lambda g(z, x) + f(z, x)]x \leq [\vartheta(z) + (1 + \lambda)\varepsilon]|x|^p + c_5|x|^r \text{ for almost all } z \in \Omega \text{ and all } x \in \mathbf{R}. \quad (3.1)$$

This unilateral growth restriction on the reaction of problem (P_λ) leads to the following auxiliary parametric nonlinear Robin problem

$$\left\{ \begin{array}{l} -\Delta_p u(z) = (\vartheta(z) + (1 + \lambda)\varepsilon)|u(z)|^{p-2}u(z) + c_5|u(z)|^{r-2}u(z) \text{ in } \Omega, \\ \frac{\partial u}{\partial n_p} + \beta(z)|u|^{p-2}u = 0 \text{ on } \partial\Omega. \end{array} \right\} \quad (7_\lambda)$$

Proposition 3.1. *If hypothesis $H(\beta)$ holds and $\lambda > 0$, then for every sufficiently small $\varepsilon > 0$ problem (7_λ) admits a positive solution*

$$\tilde{u}_\lambda \in D_+.$$

Moreover, since (7_λ) is odd, $\tilde{v}_\lambda = -\tilde{u}_\lambda \in -D_+$ is a negative solution of problem (7_λ) .

Proof. Let $\Psi_\lambda^+ : W^{1,p}(\Omega) \rightarrow \mathbf{R}$ be the C^1 -functional defined by

$$\Psi_\lambda^+(u) = \frac{1}{p}\tau(u) - \frac{1}{p} \int_\Omega [\vartheta(z) + (1 + \lambda)\varepsilon](u^+)^p dz - \frac{c_5}{r} \|u^+\|_r^r \quad \text{for all } u \in W^{1,p}(\Omega).$$

We have

$$\begin{aligned} \Psi_\lambda^+(u) &\geq c_6 \|u^-\|^p + \frac{1}{p} \left(\tau(u^+) - \int_\Omega \vartheta(z)(u^+)^p dz \right) - \frac{(1 + \lambda)\varepsilon}{p} \|u^+\|^p - c_7 \|u\|^r \\ &\quad \text{for some } c_6, c_7 > 0 \text{ (see (2.1))} \\ &\geq c_6 \|u^-\|^p + \frac{1}{p} [c_8 - (1 + \lambda)\varepsilon] \|u^+\|^p - c_7 \|u\|^r \quad \text{for all } u \in W^{1,p}(\Omega), \text{ some } c_8 > 0. \end{aligned}$$

Choosing $\varepsilon \in \left(0, \frac{c_8}{1 + \lambda}\right)$, we consider that

$$\Psi_\lambda^+(u) \geq c_9 \|u\|^p - c_7 \|u\|^r \quad \text{for all } u \in W^{1,p}(\Omega), \text{ some } c_9 > 0. \quad (3.3)$$

Since $r > p$, it follows from (3.3) that

$$u = 0 \text{ is a local minimizer of } \Psi_\lambda^+.$$

Then we can find small $\rho \in (0, 1)$ such that

$$\Psi_\lambda^+(0) = 0 < \inf \{ \Psi_\lambda^+(u) : \|u\| = \rho \} = m_\lambda^+ \text{ (see Aizicovici, Papageorgiou \& Staicu [1], proof of Proposition 29).}$$

For $t > 0$, we have

$$\begin{aligned} \Psi_\lambda^+(t\hat{u}_1) &= \frac{t^p}{p} \tau(\hat{u}_1) - \frac{t^p}{p} \int_\Omega [\vartheta(z) + (1 + \lambda)\epsilon] \hat{u}_1^p dz - \frac{t^r}{r} \|\hat{u}_1\|_r^r \\ &\leq \frac{t^p}{p} \int_\Omega [\hat{\lambda}_1 - \vartheta(z)] \hat{u}_1^p dz - \frac{t^r}{r} \|\hat{u}_1\|_r^r \\ &\leq c_{10} t^p - c_{11} t^r \text{ for some } c_{10}, c_{11} > 0. \end{aligned} \tag{3.4}$$

However, $r > p$. So, from (3.4) we have

$$\Psi_\lambda^+(t\hat{u}_1) \rightarrow -\infty \text{ as } t \rightarrow +\infty. \tag{3.5}$$

Let $k_\lambda(z, x)$ be the Carathéodory function defined by

$$k_\lambda(z, x) = [\vartheta(z) + (1 + \lambda)\epsilon] |x|^{p-2} x + c_5 |x|^{r-2} x.$$

We set

$$K_\lambda(z, x) = \int_0^x k_\lambda(z, s) ds = \frac{1}{p} [\vartheta(z) + (1 + \lambda)\epsilon] |x|^p + \frac{c_5}{r} |x|^r.$$

Recall that $p < r$ and let $q \in (p, r)$. For sufficiently big $M > 0$ we have

$$\begin{aligned} 0 < qK_\lambda(z, x) &\leq k_\lambda(z, x)x \text{ for almost all } z \in \Omega \text{ and all } |x| \geq M, \\ &\Rightarrow k_\lambda(z, \cdot) \text{ satisfies the Ambrosetti–Rabinowitz condition (see [19, p. 341])} \\ &\Rightarrow \Psi_\lambda^+(\cdot) \text{ satisfies the C-condition (see [19, p. 343]).} \end{aligned} \tag{3.6}$$

Then (3.5) and (3.6) permit the use of Theorem 2.1 (the mountain pass theorem). So, we can find $\tilde{u}_\lambda \in W^{1,p}(\Omega)$ such that

$$\tilde{u}_\lambda \in K_{\Psi_\lambda^+} \quad \text{and} \quad \Psi_\lambda^+(0) = 0 < m_\lambda^+ \leq \Psi_\lambda^+(\tilde{u}_\lambda).$$

Evidently, $\tilde{u}_\lambda \neq 0$ and we have

$$\begin{aligned} \langle A(\tilde{u}_\lambda), h \rangle &+ \int_{\partial\Omega} \beta(z) |\tilde{u}_\lambda|^{p-2} \tilde{u}_\lambda h d\sigma \\ &= \int_\Omega \{ [\vartheta(z) + (1 + \lambda)\epsilon] (\tilde{u}_\lambda^+)^{p-1} + c_5 (\tilde{u}_\lambda^+)^{r-1} \} h dz \quad \text{for all } h \in W^{1,p}(\Omega). \end{aligned} \tag{3.7}$$

In (3.7) we choose $h = -\tilde{u}_\lambda^- \in W^{1,p}(\Omega)$. Then

$$\tau(\tilde{u}_\lambda^-) = 0, \quad \Rightarrow \tilde{u}_\lambda \geq 0, \quad \tilde{u}_\lambda \neq 0 \text{ (see (2.1)).}$$

Then by (3.7) we have

$$\begin{aligned} -\Delta_p \tilde{u}_\lambda(z) &= [\vartheta(z) + (1 + \lambda)\epsilon] \tilde{u}_\lambda(z)^{p-1} + c_5 \tilde{u}_\lambda(z)^{r-1} \text{ for almost all } z \in \Omega, \\ \frac{\partial \tilde{u}_\lambda}{\partial n_p} + \beta(z) \tilde{u}_\lambda^{p-1} &= 0 \text{ on } \partial\Omega \text{ (see Papageorgiou \& Rădulescu [21]).} \end{aligned} \tag{3.8}$$

By (3.8) and Proposition 2.10 of Papageorgiou & Rădulescu [23], we have

$$\tilde{u}_\lambda \in L^\infty(\Omega).$$

So, we can apply Theorem 2 of Lieberman [17] and conclude that

$$\tilde{u}_\lambda \in C_+ \setminus \{0\}.$$

It follows from (3.8) that

$$\Delta_p \tilde{u}_\lambda(z) \leq 0 \text{ for almost all } z \in \Omega, \quad \Rightarrow \tilde{u}_\lambda \in D_+ \text{ (see Gasinski \& Papageorgiou [6, p. 738]).}$$

Since problem (7_λ) is odd, we can deduce that $\tilde{v}_\lambda = -\tilde{u}_\lambda \in -D_+$ is a negative solution of problem (7_λ) . \square

Next, we produce a uniform lower bound $\hat{c} > 0$ for the solutions \tilde{u}_λ of (7_λ) for $\lambda > 0$. It follows that $-\hat{c} < 0$ is an upper bound for the negative solutions \tilde{v}_λ .

Proposition 3.2. *If hypothesis $H(\beta)$ holds, then there exists $\hat{c} > 0$ such that*

$$\hat{c} \leq \tilde{u}_\lambda(z) \text{ and } \tilde{v}_\lambda(z) \leq -\hat{c} \text{ for all } z \in \overline{\Omega}, \lambda > 0.$$

Proof. We consider the following nonlinear Robin problem

$$\left\{ \begin{array}{l} -\Delta_p u(z) = c_5 |u(z)|^{r-2} u(z) \text{ in } \Omega, \\ \frac{\partial u}{\partial n_p} + \beta(z) |u|^{p-2} u = 0 \text{ on } \partial\Omega. \end{array} \right\} \quad (3.9)$$

We first show that problem (3.9) has a positive solution. So, let $\xi : W^{1,p}(\Omega) \rightarrow \mathbf{R}$ be the C^1 -functional defined by

$$\xi(u) = \frac{1}{p} \tau(u) - \frac{c_5}{r} \|u^+\|_r^r \text{ for all } u \in W^{1,p}(\Omega).$$

Using (2.1) we have

$$\begin{aligned} \xi(u) &\geq c_{12} \|u\|^p - c_{13} \|u\|^r \text{ for some } c_{12}, c_{13} > 0, \text{ all } u \in W^{1,p}(\Omega), \\ \Rightarrow u = 0 &\text{ is an isolated local minimizer of } \xi(\cdot) \text{ (recall that } r > p). \end{aligned}$$

So, we can find $\rho \in (0, 1)$ small such that

$$\xi(0) = 0 < \inf \{ \xi(u) : \|u\| = \rho \} = m_\xi. \quad (3.10)$$

Also, if $u \in D_+$, then

$$\xi(tu) \rightarrow -\infty \text{ as } t \rightarrow +\infty \text{ (again use the fact that } r > p). \quad (3.11)$$

Finally, since the reaction $f(x) = c_5(x^+)^{p-1}$ satisfies the Ambrosetti–Rabinowitz condition on $\mathbf{R}_+ = [0, +\infty)$, we can infer that

$$\xi(\cdot) \text{ satisfies the C-condition.} \quad (3.12)$$

Then (3.10), (3.11), (3.12) permit the use of Theorem 2.1 (the mountain pass theorem) and obtain $\bar{u} \in W^{1,p}(\Omega)$ such that

$$\bar{u} \in K_\xi \text{ and } \xi(0) = 0 < m_\xi \leq \xi(\bar{u}). \quad (3.13)$$

From (3.13) we can infer that $\bar{u} \neq 0$ and

$$\xi'(\bar{u}) = 0, \quad \Rightarrow \langle A(\bar{u}), h \rangle + \int_{\partial\Omega} \beta(z)|\bar{u}|^{p-2}\bar{u}h \, d\sigma = c_5 \int_{\Omega} (\bar{u}^+)^{p-1} h \, dz \quad \text{for all } h \in W^{1,p}(\Omega).$$

We choose $h = -\bar{u}^- \in W^{1,p}(\Omega)$. Then

$$\tau(\bar{u}^-) = 0, \quad \Rightarrow \bar{u} \geq 0, \bar{u} \neq 0 \text{ (see (2.1)).}$$

So, \bar{u} is a positive solution of (3.9). As before, the nonlinear regularity theory and the nonlinear Hopf boundary point theorem (see [6, p. 738]) imply that $\bar{u} \in D_+$.

Next, we show that there is a smallest positive solution for problem (3.9). We first observe that from Papageorgiou, Rădulescu & Repovš [24] (see the proof of Proposition 7), we know that the set S_+ of positive solutions of (3.9) is downward directed (that is, if $\bar{u}_1, \bar{u}_2 \in S_+$, then we can find $\bar{u} \in S_+$ such that $\bar{u} \leq \bar{u}_1, \bar{u} \leq \bar{u}_2$). Invoking Lemma 3.10 of Hu & Papageorgiou [14, p. 178], we can find a decreasing sequence $\{\bar{u}_n\}_{n \geq 1} \subseteq S_+ \subseteq D_+$ such that

$$\inf S_+ = \inf_{n \geq 1} \bar{u}_n.$$

We have

$$\langle A(\bar{u}_n), h \rangle + \int_{\partial\Omega} \beta(z)\bar{u}_n^{p-1} h \, d\sigma = c_5 \int_{\Omega} \bar{u}_n^{r-1} h \, dz \quad \text{for all } h \in W^{1,p}(\Omega), n \in \mathbf{N}. \tag{3.14}$$

It follows from (3.14) that $\{\bar{u}_n\}_{n \geq 1} \subseteq W^{1,p}(\Omega)$ is bounded. So, we may assume that

$$\bar{u}_n \rightharpoonup \bar{u}_* \text{ in } W^{1,p}(\Omega), \quad \bar{u}_n \rightarrow \bar{u}_* \text{ in } L^r(\Omega) \text{ and } L^p(\partial\Omega). \tag{3.15}$$

Suppose that $\bar{u}_* \equiv 0$. Let $\bar{y}_n = \frac{\bar{u}_n}{\|\bar{u}_n\|}, n \in \mathbf{N}$. Then $\|\bar{y}_n\| = 1$ for all $n \in \mathbf{N}$ and so we may assume that

$$\bar{y}_n \rightharpoonup \bar{y} \text{ in } W^{1,p}(\Omega), \quad \bar{y}_n \rightarrow \bar{y} \text{ in } L^p(\Omega) \text{ and } L^p(\partial\Omega). \tag{3.16}$$

From (3.14) we have

$$\langle A(\bar{y}_n), h \rangle + \int_{\partial\Omega} \beta(z)\bar{y}_n^{p-1} h \, d\sigma = c_5 \int_{\Omega} \bar{u}_n^{r-p} \bar{y}_n^{p-1} h \, dz \quad \text{for all } h \in W^{1,p}(\Omega), n \in \mathbf{N}.$$

Choosing $h = \bar{y}_n - \bar{y} \in W^{1,p}(\Omega)$, passing to the limit as $n \rightarrow \infty$, and using (3.16) and the fact that $\bar{u}_* = 0$, we obtain

$$\lim_{n \rightarrow \infty} \langle A(\bar{y}_n), \bar{y}_n - \bar{y} \rangle = 0, \quad \Rightarrow \bar{y}_n \rightarrow \bar{y} \text{ in } W^{1,p}(\Omega), \|\bar{y}\| = 1. \text{ (see Proposition 2.4).} \tag{3.17}$$

Passing to the limit as $n \rightarrow \infty$ in (3.16), and using (3.17) and the fact that $\bar{u}_* = 0$, we obtain

$$\begin{aligned} \langle A(\bar{y}), h \rangle + \int_{\partial\Omega} \beta(z)\bar{y}^{p-1} h \, d\sigma &= 0 \quad \text{for all } h \in W^{1,p}(\Omega), \\ \Rightarrow \tau(\bar{y}) &= 0, \\ \Rightarrow \bar{y} &= 0 \text{ (see (2.1)), contradicting (3.17).} \end{aligned}$$

So, $\bar{u}_* \neq 0$. In (3.14) we choose $h = \bar{u}_n - \bar{u}_* \in W^{1,p}(\Omega)$, pass to the limit as $n \rightarrow \infty$, and use (3.15) and Proposition 2.4. Then

$$\bar{u}_n \rightarrow \bar{u}_* \text{ in } W^{1,p}(\Omega).$$

Hence, in the limit as $n \rightarrow \infty$ in (3.14), we obtain

$$\langle A(\bar{u}_*), h \rangle + \int_{\partial\Omega} \beta(z)\bar{u}_*^{p-1} h \, d\sigma = c_5 \int_{\Omega} \bar{u}_*^{r-1} h \, dz \quad \text{for all } h \in W^{1,p}(\Omega) \quad \Rightarrow \bar{u}_* \in S_+ \text{ and } \bar{u}_* = \inf S_+.$$

Now let $\tilde{u}_\lambda \in D_+$ be a solution of (7_λ) (see Proposition 3.1). We consider the Carathéodory function $\gamma(z, x)$ defined by

$$\gamma(z, x) = \begin{cases} c_5(x^+)^{r-1} & \text{if } x \leq \tilde{u}_\lambda(z), \\ c_5\tilde{u}_\lambda(z)^{r-1} & \text{if } \tilde{u}_\lambda(z) < x. \end{cases} \tag{3.18}$$

We set $\Gamma(z, x) = \int_0^x \gamma(z, s) ds$ and consider the C^1 -functional $\hat{\xi} : W^{1,p}(\Omega) \rightarrow \mathbf{R}$ defined by

$$\hat{\xi}(u) = \frac{1}{p} \tau(u) - \int_\Omega \Gamma(z, u) dz \quad \text{for all } u \in W^{1,p}(\Omega).$$

It follows by (2.1) and (3.18) that $\hat{\xi}(\cdot)$ is coercive. Also, it is sequentially weakly lower semicontinuous. So, we can find $\bar{u} \in W^{1,p}(\Omega)$ such that

$$\hat{\xi}(\bar{u}) = \inf \{ \hat{\xi}(u) : u \in W^{1,p}(\Omega) \}. \tag{3.19}$$

As before, since $r > p$, we have $\hat{\xi}(\bar{u}) < 0 = \hat{\xi}(0)$, hence $\bar{u} \neq 0$. From (3.19) we have

$$\langle A(\bar{u}), h \rangle + \int_{\partial\Omega} \beta(z)|\bar{u}|^{p-2}\bar{u}h d\sigma = \int_\Omega \gamma(z, \bar{u})h dz \quad \text{for all } h \in W^{1,p}(\Omega). \tag{3.20}$$

In (3.20) we first choose $h = -\bar{u}^- \in W^{1,p}(\Omega)$. We obtain

$$\tau(\bar{u}^-) = 0 \text{ (see (3.18)), } \Rightarrow \bar{u} \geq 0, \bar{u} \neq 0 \text{ (see (2.1)).}$$

Next, we choose $h = (\bar{u} - \tilde{u}_\lambda)^+ \in W^{1,p}(\Omega)$ in (3.20). Then

$$\begin{aligned} \langle A(\bar{u}), (\bar{u} - \tilde{u}_\lambda)^+ \rangle + \int_{\partial\Omega} \beta(z)\bar{u}^{p-1}(\bar{u} - \tilde{u}_\lambda)^+ d\sigma &= \int_\Omega c_5\tilde{u}_\lambda^{r-1}(\bar{u} - \tilde{u}_\lambda)^+ dz \\ &\leq \int_\Omega ([\vartheta(z) + (1 + \lambda)\varepsilon]\tilde{u}_\lambda^{p-1} + c_5\tilde{u}_\lambda^{r-1})(\bar{u} - \tilde{u}_\lambda)^+ dz \\ &= \langle A(\tilde{u}_\lambda), (\bar{u} - \tilde{u}_\lambda)^+ \rangle + \int_{\partial\Omega} \beta(z)\tilde{u}_\lambda^{p-1}(\bar{u} - \tilde{u}_\lambda)^+ d\sigma \\ &\Rightarrow \bar{u} \leq \tilde{u}_\lambda \text{ (by Proposition 2.4).} \end{aligned}$$

So, we have proved that

$$\bar{u} \in [0, \tilde{u}_\lambda], \quad \bar{u} \neq 0.$$

It follows by (3.18), (3.20) and the above relation that $\bar{u} \in S_+ \subseteq D_+$. Therefore

$$0 < \hat{c} = \min_\Omega \bar{u}_* \leq \tilde{u}_\lambda \quad \text{for all } \lambda > 0.$$

The oddness of (3.9) implies that $\tilde{v}_\lambda \leq -\hat{c} < 0$ for all $\lambda > 0$. □

Now we are ready to produce two nontrivial constant sign solutions when $\lambda > 0$ is big enough.

Proposition 3.3. *If hypotheses $H(\beta)$, $H(g)$, $H(f)$, H_0 hold, then for sufficiently large $\lambda > 0$ problem (P_λ) has two constant sign solutions*

$$u_0 \in \text{int}_{C^1(\bar{\Omega})} [0, \tilde{u}_\lambda], \quad v_0 \in \text{int}_{C^1(\bar{\Omega})} [\tilde{v}_\lambda, 0],$$

with $\tilde{u}_\lambda \in D_+$ and $\tilde{v}_\lambda \in -D_+$ constant sign solutions of (7_λ) .

Proof. We introduce the following truncation of the reaction in problem (P_λ) :

$$\eta_\lambda^+(z, x) = \begin{cases} \lambda g(z, x^+) + f(z, x^+) & \text{if } x \leq \tilde{u}_\lambda(z), \\ \lambda g(z, \tilde{u}_\lambda(z)) + f(z, \tilde{u}_\lambda(z)) & \text{if } \tilde{u}_\lambda(z) < x. \end{cases} \tag{3.21}$$

This is a Carathéodory function. We set $H_\lambda^+(z, x) = \int_0^x \eta_\lambda^+(z, s) ds$ and consider the C^1 -functional $d_\lambda^+ : W^{1,p}(\Omega) \rightarrow \mathbf{R}$ defined by

$$d_\lambda^+(u) = \frac{1}{p} \tau(u) - \int_\Omega H_\lambda^+(z, u) dz \quad \text{for all } u \in W^{1,p}(\Omega).$$

From (2.1) and (3.21) we see that $d_\lambda^+(\cdot)$ is coercive. Also, it is sequentially weakly lower semicontinuous. So, we can find $u_0 \in W^{1,p}(\Omega)$ such that

$$d_\lambda^+(u_0) = \inf \{ d_\lambda^+(u) : u \in W^{1,p}(\Omega) \}. \tag{3.22}$$

Let $c \in (0, \hat{c})$ with $\hat{c} > 0$ as in Proposition 3.2. Then for all $\lambda > 0$, we have

$$d_\lambda^+(c) = \frac{c^p}{p} \int_{\partial\Omega} \beta(z) d\sigma - \lambda \int_\Omega G(z, c) dz - \int_\Omega F(z, c) dz.$$

Note that $\int_\Omega F(z, c) dz > 0$ (see hypothesis $H(g)(i)$). So,

$$\begin{aligned} d_\lambda^+(c) &< 0 \text{ for sufficiently large } \lambda > 0, \\ \Rightarrow d_\lambda^+(u_0) &< 0 = d_\lambda^+(0) \text{ for sufficiently large } \lambda > 0, \\ \Rightarrow u_0 &\neq 0. \end{aligned}$$

From (3.22) we have

$$\langle A(u_0), h \rangle + \int_{\partial\Omega} \beta(z) |u_0|^{p-2} u_0 h d\sigma = \int_\Omega \eta_\lambda^+(z, u_0) h dz \quad \text{for all } h \in W^{1,p}(\Omega). \tag{3.23}$$

In (3.23) we first choose $h = -u_0^- \in W^{1,p}(\Omega)$ and obtain

$$u_0 \geq 0, \quad u_0 \neq 0.$$

Then in (3.23) we choose $h = (u_0 - \tilde{u}_\lambda)^+ \in W^{1,p}(\Omega)$. As in the proof of Proposition 3.2, using this time (3.1), we obtain

$$u_0 \leq \tilde{u}_\lambda.$$

So, we have proved that

$$u_0 \in [0, \tilde{u}_\lambda], \quad u_0 \neq 0. \tag{3.24}$$

By (3.21), (3.23), (3.24) and Theorem 2 of Lieberman [17], we have

$$u_0 \in C_+ \setminus \{0\} \text{ is a positive solution of } (P_\lambda), \quad \lambda > 0 \text{ big.}$$

Therefore we have

$$\Delta_p u_0(z) \leq 0 \text{ for almost all } z \in \Omega, \quad \Rightarrow u_0 \in D_+ \text{ (see Gasinski \& Papageorgiou [6, p. 738]).}$$

Also, we have

$$\begin{aligned} -\Delta_p u_0(z) &= \lambda g(z, u_0(z)) + f(z, u_0(z)) \\ &\leq \lambda g(z, \tilde{u}_\lambda(z)) + f(z, \tilde{u}_\lambda(z)) \text{ (see (3.24) and hypothesis } H_0) \\ &\leq [\vartheta(z) + (1 + \lambda)\varepsilon] \tilde{u}_\lambda(z)^{p-1} + c_5 \tilde{u}_\lambda(z)^{r-1} \text{ (see (3.1))} \\ &= -\Delta_p \tilde{u}_\lambda(z) \text{ for almost all } z \in \Omega, \\ \Rightarrow \tilde{u}_\lambda - u_0 &\in \text{int } \hat{C}_+ \text{ (see Proposition 2.3).} \end{aligned}$$

We conclude that

$$u_0 \in \text{int}_{C^1(\bar{\Omega})} [0, \tilde{u}_\lambda].$$

For the negative solution, we introduce the Carathéodory function $\eta_\lambda^-(z, x)$ defined by

$$\eta_\lambda^-(z, x) = \begin{cases} \lambda g(z, \tilde{v}_\lambda(z)) + f(z, \tilde{v}_\lambda(z)) & \text{if } x < \tilde{v}_\lambda(z), \\ \lambda g(z, -x^-) + f(z, -x^-) & \text{if } \tilde{v}_\lambda(z) \leq x. \end{cases}$$

We set $H_\lambda^-(z, x) = \int_0^x \eta_\lambda^-(z, s) ds$ and consider the C^1 -functional $d_\lambda^- : W^{1,p}(\Omega) \rightarrow \mathbf{R}$ defined by

$$d_\lambda^-(u) = \frac{1}{p} \tau(u) - \int_\Omega H_\lambda^-(z, u) dz \quad \text{for all } u \in W^{1,p}(\Omega).$$

Working as above, this time with the functional $d_\lambda^-(\cdot)$, we produce a solution v_0 of (P_λ) for large enough $\lambda > 0$ such that

$$v_0 \in -D_+, \quad v_0 \in \text{int}_{C^1(\bar{\Omega})} [\tilde{v}_\lambda, 0].$$

The proof is now complete. □

Using $u_0 \in D_+$ and $v_0 \in -D_+$ from Proposition 3.3, we will produce two more constant sign solutions.

Proposition 3.4. *If hypotheses $H(\beta)$, $H(g)$, $H(f)$, H_0 hold, then for $\lambda > 0$ big enough, problem (P_λ) admits two more constant sign solutions $\hat{u} \in D_+$ and $\hat{v} \in -D_+$ such that*

$$\hat{u} - u_0 \in \text{int } C_+, \quad v_0 - \hat{v} \in \text{int } C_+$$

with $u_0 \in D_+$ and $v_0 \in -D_+$ the solutions from Proposition 3.3.

Proof. We introduce the following truncation of the reaction in problem (P_λ) :

$$i_\lambda^+(z, x) = \begin{cases} \lambda g(z, u_0(z)) + f(z, u_0(z)) & \text{if } x \leq u_0(z), \\ \lambda g(z, x) + f(z, x) & \text{if } u_0(z) < x. \end{cases} \tag{3.25}$$

This is a Carathéodory function. We set $I_\lambda^+(z, x) = \int_0^x i_\lambda^+(z, s) ds$ and consider the C^1 -functional $\chi_\lambda^+ : W^{1,p}(\Omega) \rightarrow \mathbf{R}$ defined by

$$\chi_\lambda^+(u) = \frac{1}{p} \tau(u) - \int_\Omega I_\lambda^+(z, u) dz \quad \text{for all } u \in W^{1,p}(\Omega).$$

Claim 3.5. $\chi_\lambda^+(\cdot)$ satisfies the C -condition.

We consider a sequence $\{u_n\}_{n \geq 1} \subseteq W^{1,p}(\Omega)$ such that

$$\left| \chi_\lambda^+(u_n) \right| \leq M_1 \text{ for some } M_1 > 0 \text{ and all } n \in \mathbf{N}, \tag{3.26}$$

$$(1 + \|u_n\|)(\chi_\lambda^+)'(u_n) \rightarrow 0 \text{ in } W^{1,p}(\Omega)^* \text{ as } n \rightarrow \infty. \tag{3.27}$$

From (3.27) we have

$$\left| \langle A(u_n), h \rangle + \int_{\partial\Omega} \beta(z) |u_n|^{p-2} u_n h d\sigma - \int_\Omega i_\lambda^+(z, u_n) h dz \right| \leq \frac{\epsilon_n \|h\|}{1 + \|u_n\|} \quad \text{for all } h \in W^{1,p}(\Omega), \text{ with } \epsilon_n \rightarrow 0^+. \tag{3.28}$$

In (3.28) we choose $h = -u_n^- \in W^{1,p}(\Omega)$. Then

$$\begin{aligned} \tau(u_n^-) &\leq c_{14} \|u_n^-\| \text{ for some } c_{14} > 0 \text{ and all } n \in \mathbf{N} \quad (\text{see (3.25) and hypotheses } H(g)(i), H(f)(i)), \\ &\Rightarrow \{u_n^-\}_{n \geq 1} \subseteq W^{1,p}(\Omega) \text{ is bounded (see (2.2)).} \end{aligned} \tag{3.29}$$

Using (3.29) in (3.28), we obtain

$$\left| \langle A(u_n^+), h \rangle + \int_{\partial\Omega} \beta(z)(u_n^+)^{p-1} h \, d\sigma - \int_{\Omega} i_\lambda^+(z, u_n^+) \, dz \right| \leq c_{15} \|h\| \tag{3.30}$$

for some $c_{15} > 0$ and all $h \in W^{1,p}(\Omega)$, $n \in \mathbf{N}$.

We will show that $\{u_n^+\}_{n \geq 1} \subseteq W^{1,p}(\Omega)$ is bounded, too. Arguing by contradiction, suppose that $\|u_n^+\| \rightarrow \infty$ as $n \rightarrow \infty$. Let $y_n = \frac{u_n^+}{\|u_n^+\|}$, $n \in \mathbf{N}$. Then $\|y_n\| = 1$, $y_n \geq 0$ for all $n \in \mathbf{N}$ and so we may assume that

$$y_n \xrightarrow{w} y \text{ in } W^{1,p}(\Omega) \quad \text{and} \quad y_n \rightarrow y \text{ in } L^p(\Omega) \text{ and } L^p(\partial\Omega), \quad y \geq 0. \tag{3.31}$$

From (3.30) we have

$$\left| \langle A(y_n), h \rangle + \int_{\partial\Omega} \beta(z)y_n^{p-1} h \, d\sigma - \int_{\Omega} \frac{i_\lambda^+(z, u_n^+)}{\|u_n^+\|^{p-1}} h \, dz \right| \leq \frac{c_{15} \|h\|}{\|u_n^+\|^{p-1}} \quad \text{for all } n \in \mathbf{N}, h \in W^{1,p}(\Omega). \tag{3.32}$$

From (2.5) and (3.25), we see that

$$\left\{ \frac{i_\lambda^+(\cdot, u_n^+(\cdot))}{\|u_n^+\|^{p-1}} \right\}_{n \geq 1} \subseteq L^{p'}(\Omega) \text{ is bounded } \left(\frac{1}{p} + \frac{1}{p'} = 1 \right). \tag{3.33}$$

Passing to a subsequence if necessary, and using hypotheses $H(g)(ii)$, $H(f)(ii)$, we have

$$\frac{i_\lambda^+(\cdot, u_n^+(\cdot))}{\|u_n^+\|^{p-1}} \xrightarrow{w} \hat{\lambda}_m y^{p-1} \text{ in } L^{p'}(\Omega) \tag{3.34}$$

(see Aizicovici, Papageorgiou & Staicu [1], proof of Proposition 4.8).

In (3.32) we choose $h = y_n - y \in W^{1,p}(\Omega)$, pass to the limit as $n \rightarrow \infty$, and use (3.31) and (3.33). Then

$$\lim_{n \rightarrow \infty} \langle A(y_n), y_n - y \rangle = 0, \quad \Rightarrow y_n \rightarrow y \text{ in } W^{1,p}(\Omega) \text{ (see Proposition 2.4) and so } \|y\| = 1, y \geq 0. \tag{3.35}$$

So, if in (3.32) we pass to the limit as $n \rightarrow \infty$, and use (3.34) and (3.35) to obtain

$$\begin{aligned} \langle A(y), h \rangle + \int_{\partial\Omega} \beta(z)y^{p-1} h \, d\sigma &= \hat{\lambda}_m \int_{\Omega} y^{p-1} h \, dz \quad \text{for all } h \in W^{1,p}(\Omega), \\ \Rightarrow -\Delta_p y(z) &= \hat{\lambda}_m y(z)^{p-1} \text{ for almost all } z \in \Omega, \quad \frac{\partial y}{\partial n_p} + \beta(z)y^{p-1} = 0 \text{ on } \partial\Omega \end{aligned} \tag{3.36}$$

(see Papageorgiou & Rădulescu [21]).

Since $m \geq 2$, it follows by (3.36) that $y(\cdot)$ must be nodal, a contradiction to (3.35). Therefore

$$\{u_n^+\}_{n \geq 1} \subseteq W^{1,p}(\Omega) \text{ is bounded, } \Rightarrow \{u_n\}_{n \geq 1} \subseteq W^{1,p}(\Omega) \text{ is bounded (see (3.29)).}$$

So, we may assume that

$$u_n \xrightarrow{w} u \text{ in } W^{1,p}(\Omega) \quad \text{and} \quad u_n \rightarrow u \text{ in } L^p(\Omega) \text{ and in } L^p(\partial\Omega). \tag{3.37}$$

In (3.28) we choose $h = u_n - u \in W^{1,p}(\Omega)$, pass to the limit as $n \rightarrow \infty$, and use (3.37) and the fact that $\{i_\lambda^+(\cdot, u_n(\cdot))\}_{n \geq 1} \subseteq L^{p'}(\Omega)$ is bounded (see (2.5) and (3.25)). We obtain

$$\lim_{n \rightarrow \infty} \langle A(u_n), u_n - u \rangle = 0, \quad \Rightarrow u_n \rightarrow u \text{ in } W^{1,p}(\Omega) \text{ (see Proposition 2.4).}$$

So $\mathcal{I}_\lambda^+(\cdot)$ satisfies the C -condition. This proves Claim 3.5.

Claim 3.6. We may assume that $u_0 \in D_+$ is a local minimizer of $\chi_\lambda^+(\cdot)$.

For sufficiently large $\lambda > 0$, as in Proposition 3.3, let $\tilde{u}_\lambda \in D_+$ be a solution of (7_λ) (see Proposition 3.1). From Proposition 3.3 we know that

$$\tilde{u}_\lambda - u_0 \in \text{int } C_+. \quad (3.38)$$

We introduce the following truncation of $i_\lambda^+(z, \cdot)$:

$$j_\lambda^+(z, x) = \begin{cases} i_\lambda^+(z, x) & \text{if } x \leq \tilde{u}_\lambda(z) \\ i_\lambda^+(z, \tilde{u}_\lambda(z)) & \text{if } \tilde{u}_\lambda(z) < x. \end{cases} \quad (3.39)$$

This is a Carathéodory function. We set $J_\lambda^+(z, x) = \int_0^x j_\lambda^+(z, s) ds$ and consider the C^1 -functional $\hat{\chi}_\lambda^+ : W^{1,p}(\Omega) \rightarrow \mathbf{R}$ defined by

$$\hat{\chi}_\lambda^+(u) = \frac{1}{p} \tau(u) - \int_\Omega J_\lambda^+(z, u) dz \quad \text{for all } u \in W^{1,p}(\Omega).$$

By (2.1) and (3.39), it is clear that $\hat{\chi}_\lambda^+(\cdot)$ is coercive. Also, it is sequentially weakly lower semicontinuous. Hence we can find $\hat{u}_0 \in W^{1,p}(\Omega)$ such that

$$\hat{\chi}_\lambda^+(\hat{u}_0) = \inf \{ \hat{\chi}_\lambda^+(u) : u \in W^{1,p}(\Omega) \}. \quad (3.40)$$

Using (3.39), the nonlinear regularity theory and the nonlinear maximum principle, we can easily show that

$$K_{\hat{\chi}_\lambda^+} \subseteq [u_0, \tilde{u}_\lambda] \cap D_+. \quad (3.41)$$

Evidently, $\tilde{u}_\lambda \notin K_{\hat{\chi}_\lambda^+}$ (see (3.1)). So, from (3.40) and (3.41), we have

$$\hat{u}_0 \in [u_0, \tilde{u}_\lambda] \cap D_+, \quad \hat{u}_0 \neq \tilde{u}_\lambda.$$

If $\hat{u}_0 \neq u_0$, then this is the desired second positive solution of (P_λ) for sufficiently large $\lambda > 0$, and using Proposition 2.3, we have

$$\hat{u}_0 - u_0 \in \text{int } C_+.$$

Therefore we are done.

So, we may assume that $\hat{u}_0 = u_0 \in D_+$. Note that

$$\chi_\lambda^+ \Big|_{[0, \tilde{u}_\lambda]} = \hat{\chi}_\lambda^+ \Big|_{[0, \tilde{u}_\lambda]} \quad (\text{see (3.39)}). \quad (3.42)$$

From Proposition 3.3, we have

$$u_0 \in \text{int}_{C^1(\overline{\Omega})} [0, \tilde{u}_\lambda]. \quad (3.43)$$

Then it follows from (3.40), (3.41), (3.42) that

$$u_0 \text{ is a local } C^1(\overline{\Omega})\text{-minimizer of } \chi_\lambda^+(\cdot), \quad \Rightarrow \quad u_0 \text{ is a local } W^{1,p}(\Omega)\text{-minimizer of } \chi_\lambda^+(\cdot) \quad (\text{see Proposition 2.2}).$$

This proves Claim 3.6.

Using (3.25), we can show that

$$K_{\chi_\lambda^+} \subseteq [u_0] \cap D_+.$$

So, we may assume that $K_{\chi_\lambda^+}$ is finite, or otherwise we already have an infinity of positive solutions of (P_λ) (for large enough $\lambda > 0$) strictly bigger than u_0 and so we are done. Then on account of Claim 3.6, we can find sufficiently small $\rho \in (0, 1)$ such that

$$\chi_\lambda^+(u_0) < \inf \{ \chi_\lambda^+(u) : \|u - u_0\| = \rho \} = \tilde{m}_\lambda^+ \text{ (see [1]).} \tag{3.44}$$

From hypotheses $H(g)(ii)$, $H(f)(ii)$ and since $m \geq 2$, we have

$$\chi_\lambda^+(t\hat{u}_1) \rightarrow -\infty \text{ as } t \rightarrow +\infty. \tag{3.45}$$

Then (3.44), (3.45) and Claim 3.5 permit the use of Theorem 2.1 (the mountain pass theorem). So, we can find $\hat{u} \in W^{1,p}(\Omega)$ such that

$$\hat{u} \in K_{\chi_\lambda^+} \subseteq [u_0] \cap D_+ \quad \text{and} \quad \tilde{m}_\lambda^+ \leq \chi_\lambda^+(\hat{u}). \tag{3.46}$$

It follows from (3.44), (3.46) and (3.25) that

$$\hat{u} \in D_+ \text{ is a second positive solution of } (P_\lambda) \text{ for sufficiently large } \lambda > 0, \quad u_0 \leq \hat{u}, \quad u_0 \neq \hat{u}.$$

We have

$$\begin{aligned} -\Delta_p u_0(z) &= \lambda g(z, u_0(z)) + f(z, u_0(z)) \\ &\leq \lambda g(z, \hat{u}(z)) + f(z, \hat{u}(z)) \text{ (see (3.46) and hypothesis } H_0) \\ &= -\Delta_p \hat{u}(z) \text{ for almost all } z \in \Omega. \end{aligned} \tag{3.47}$$

Note that $\lambda g(\cdot, u_0(\cdot)) + f(\cdot, u_0(\cdot)) \neq \lambda g(\cdot, \hat{u}(\cdot)) + f(\cdot, \hat{u}(\cdot))$ (see hypothesis H_0). So, from (3.47) and Proposition 2.3, we can infer that

$$\hat{u} - u_0 \in \text{int } \hat{C}_+.$$

Similarly, for the second negative solution, we use $v_0 \in -D_+$ from Proposition 3.3. So, we define

$$i_\lambda^-(z, x) = \begin{cases} \lambda g(z, x) + f(z, x) & \text{if } x \leq v_0(z), \\ \lambda g(z, v_0(z)) + f(z, v_0(z)) & \text{if } v_0(z) < x. \end{cases}$$

This is a Carathéodory function. We set $I_\lambda^-(z, x) = \int_0^x i_\lambda^-(z, s) ds$ and consider the C^1 -functional $\chi_\lambda^- : W^{1,p}(\Omega) \rightarrow \mathbf{R}$ defined by

$$\chi_\lambda^-(u) = \frac{1}{p} \tau(u) - \int_\Omega I_\lambda^-(z, u) dz \quad \text{for all } u \in W^{1,p}(\Omega).$$

Working as above, this time with χ_λ^- and truncating at $\tilde{v}_\lambda \in -D_+$ to produce $\hat{\chi}_\lambda^-(\cdot)$, we generate a second negative solution \hat{v} of problem (P_λ) for sufficiently large $\lambda > 0$, such that

$$\hat{v} \in -D_+ \quad \text{and} \quad v_0 - \hat{v} \in \text{int } \hat{C}_+.$$

This completes the proof. □

4 | THE FIFTH SOLUTION

So far we have four nontrivial smooth solutions, all with sign information (two positive and two negative). In this section, using the theory of critical groups, we establish the existence of a fifth nontrivial smooth solution distinct from the other four.

Proposition 4.1. *If hypotheses $H(\beta)$, $H(g)$, $H(f)$, H_0 hold and $\lambda > 0$ is sufficiently large, then $C_k(\chi_\lambda^\pm, \infty) = 0$ for all $k \in \mathbf{N}_0$.*

Proof. We present the proof for the functional $\chi_\lambda^+(\cdot)$, the proof for $\chi_\lambda^-(\cdot)$ being similar.

Let $\lambda \in (\hat{\lambda}_m, \hat{\lambda}_{m+1}) \setminus \hat{\sigma}(p)$ (recall that $\hat{\sigma}(p)$ denotes the set of eigenvalues of $-\Delta_p$ with Robin boundary condition) and consider the C^1 -functional $\psi_\lambda^+ : W^{1,p}(\Omega) \rightarrow \mathbf{R}$ defined by

$$\psi_\lambda^+(u) = \frac{1}{p} \tau(u) - \frac{\lambda}{p} \|u^+\|_p^p \quad \text{for all } u \in W^{1,p}(\Omega).$$

We consider the following homotopy

$$h_\lambda^+(t, u) = (1-t)\chi_\lambda^+(u) + t\psi_\lambda^+(u) \quad \text{for all } (t, u) \in [0, 1] \times W^{1,p}(\Omega).$$

Claim 4.2. We can find $\eta_0 \in \mathbf{R}$ and $\delta_0 > 0$ such that

$$h_\lambda^+(t, u_0) \leq \eta_0 \Rightarrow (1 + \|u\|) \left\| (h_\lambda^+)'_u(t, u) \right\|_* \geq \delta_0 \quad \text{for all } t \in [0, 1].$$

To prove Claim 4.2, we argue indirectly. So, suppose Claim 4.2 is not true. Evidently, $h_\lambda^+(\cdot, \cdot)$ maps bounded sets to bounded ones. Hence we can find $\{t_n\}_{n \geq 1} \subseteq [0, 1]$ and $\{u_n\}_{n \geq 1} \subseteq W^{1,p}(\Omega)$ such that

$$t_n \rightarrow t, \|u_n\| \rightarrow \infty, h_\lambda^+(t_n, u_n) \rightarrow -\infty \quad \text{and} \quad (1 + \|u_n\|) (h_\lambda^+)'_{t_n}(t_n, u_n) \rightarrow 0. \quad (4.1)$$

From the last convergence in (4.1), we have

$$\begin{aligned} & \left| \langle A(u_n), h \rangle + \int_{\partial\Omega} \beta(z) |u_n|^{p-2} u_n h \, d\sigma - (1-t_n) \int_{\Omega} i_\lambda^+(z, u_n) h \, dz - \lambda t_n \int_{\Omega} (u_n^+)^{p-1} h \, dz \right| \\ & \leq \frac{\epsilon_n \|h\|}{1 + \|u_n\|} \quad \text{for all } h \in W^{1,p}(\Omega), \text{ with } \epsilon_n \rightarrow 0^+. \end{aligned} \quad (4.2)$$

In (4.2) we choose $h = -u_n^- \in W^{1,p}(\Omega)$. Then

$$\tau(u_n^-) \leq c_{16} \|u_n^-\| \quad \text{for some } c_{16} > 0 \text{ and all } n \in \mathbf{N} \text{ (see (3.25)),} \quad \Rightarrow \{u_n^-\}_{n \geq 1} \subseteq W^{1,p}(\Omega) \text{ is bounded (see (2.1)).} \quad (4.3)$$

From (4.1) and (4.3) it follows that

$$\|u_n^+\| \rightarrow +\infty.$$

We set $y_n = \frac{u_n^+}{\|u_n^+\|}$, $n \in \mathbf{N}$. Then $\|y_n\| = 1$, $y_n \geq 0$ for all $n \in \mathbf{N}$. So, we may assume that

$$y_n \xrightarrow{w} y \text{ in } W^{1,p}(\Omega) \quad \text{and} \quad y_n \rightarrow y \text{ in } L^p(\Omega) \text{ and in } L^p(\partial\Omega), \quad y \geq 0. \quad (4.4)$$

From (4.2) and (4.3) it follows that

$$\begin{aligned} & \left| \langle A(y_n), h \rangle + \int_{\partial\Omega} \beta(z) y_n^{p-1} h \, d\sigma - (1-t_n) \int_{\Omega} \frac{i_\lambda^+(z, u_n^+)}{\|u_n^+\|^{p-1}} h \, dz - \lambda t_n \int_{\Omega} y_n^{p-1} h \, dz \right| \leq \epsilon'_n \|h\| \\ & \text{for all } h \in W^{1,p}(\Omega), \text{ with } \epsilon'_n \rightarrow 0^+. \end{aligned} \quad (4.5)$$

From (2.5) and (3.25), we see that

$$\left\{ \frac{i_\lambda^+(\cdot, u_n^+(\cdot))}{\|u_n^+\|^{p-1}} \right\}_{n \geq 1} \subseteq L^{p'}(\Omega) \text{ is bounded.} \quad (4.6)$$

Passing to a subsequence if necessary and using hypotheses $H(g)(ii)$ and $H(f)(ii)$ we have

$$\frac{i_\lambda^+(\cdot, u_n^+(\cdot))}{\|u_n^+\|^{p-1}} \rightharpoonup \hat{\lambda}_m y^{p-1} \text{ in } L^{p'}(\Omega) \text{ (see [1]).} \tag{4.7}$$

In (4.5) we choose $h = y_n - y \in W^{1,p}(\Omega)$, pass to the limit as $n \rightarrow \infty$, and use (4.4), (4.6). Then

$$\lim_{n \rightarrow \infty} \langle A(y_n), y_n - y \rangle = 0, \Rightarrow y_n \rightarrow y \text{ in } W^{1,p}(\Omega) \text{ (see Proposition 2.4), hence } \|y\| = 1, y \geq 0. \tag{4.8}$$

In (4.5) we pass to the limit as $n \rightarrow \infty$, and use (4.7), (4.8) and the continuity of $A(\cdot)$ (Proposition 2.4). We obtain

$$\begin{aligned} \langle A(y), h \rangle + \int_{\partial\Omega} \beta(z) y^{p-1} h \, d\sigma &= [(1-t)\hat{\lambda}_m + t\lambda] \int_{\Omega} y^{p-1} h \, dz \text{ for all } h \in W^{1,p}(\Omega), \\ \Rightarrow -\Delta_p y(z) &= \lambda_t y(z)^{p-1} \text{ for almost all } z \in \Omega, \frac{\partial y}{\partial n_p} + \beta(z) y^{p-1} = 0 \text{ on } \partial\Omega, \end{aligned} \tag{4.9}$$

with $\lambda_t = (1-t)\hat{\lambda}_m + t\lambda$. We have

$$\lambda_t \in [\hat{\lambda}_m, \hat{\lambda}_{m+1}). \tag{4.10}$$

From (4.9) and (4.10) and since $m \geq 2$, we can infer that

$$y = 0 \text{ or } y \text{ is nodal.}$$

Both assertions contradict (4.8). This proves Claim 4.2.

Then Claim 4.2 and Theorem 5.1.21 of Chang [3, p. 334] (see also Liang & Su [16, Proposition 3.2]), imply that

$$C_k(h_\lambda^+(0, \cdot, \infty)) = C_k(h_\lambda^+(1, \cdot, \infty)) \text{ for all } k \in \mathbf{N}_0, \Rightarrow C_k(\chi_\lambda^+, \infty) = C_k(\psi_\lambda^+, \infty) \text{ for all } k \in \mathbf{N}_0. \tag{4.11}$$

Now we consider the following homotopy

$$\hat{h}_\lambda^+(t, u) = \psi_\lambda^+(u) - t \int_{\Omega} u \, dz \text{ for all } (t, u) \in [0, 1] \times W^{1,p}(\Omega).$$

Claim 4.3. $(\hat{h}_\lambda^+)'_u(t, u) \neq 0$ for all $t \in [0, 1], u \in W^{1,p}(\Omega) \setminus \{0\}$.

Again, we argue indirectly. So, suppose that for some $t \in [0, 1]$ and $u \in W^{1,p}(\Omega) \setminus \{0\}$, we have

$$\begin{aligned} (\hat{h}_\lambda^+)'_u(t, u) &= 0, \\ \Rightarrow \langle A(u), h \rangle + \int_{\partial\Omega} \beta(z) |u|^{p-2} u h \, d\sigma &= \lambda \int_{\Omega} (u^+)^{p-1} h \, dz + t \int_{\Omega} h \, dz \text{ for all } h \in W^{1,p}(\Omega). \end{aligned} \tag{4.12}$$

In (4.12) we choose $h = -u^- \in W^{1,p}(\Omega)$. Then

$$\tau(u^-) \leq 0, \Rightarrow u \geq 0, u \neq 0 \text{ (see (2.1)).}$$

Hence (4.12) becomes

$$\begin{aligned} \langle A(u), h \rangle + \int_{\partial\Omega} \beta(z) u^{p-1} h \, dz &= \lambda \int_{\Omega} u^{p-1} h \, dz + t \int_{\Omega} h \, dz \text{ for all } h \in W^{1,p}(\Omega), \\ \Rightarrow -\Delta_p u(z) &= \lambda u(z)^{p-1} + t \text{ for almost all } z \in \Omega, \frac{\partial u}{\partial n_p} + \beta(z) u^{p-1} = 0 \text{ on } \partial\Omega. \end{aligned} \tag{4.13}$$

As before, the nonlinear regularity theory implies that $u \in C_+ \setminus \{0\}$. Also, from (4.13) we have

$$\Delta_p u(z) \leq 0 \text{ for almost all } z \in \Omega, \Rightarrow u \in D_+ \text{ (see Gasinski \& Papageorgiou [6, p. 738]).}$$

Let $v \in D_+$ and consider the function $R(v, u)(\cdot)$ from Section 2. Using Proposition 2.6, we get

$$\begin{aligned} 0 &\leq \int_{\Omega} R(v, u) \, dz \\ &= \|Dv\|_p^p - \int_{\Omega} (-\Delta_p u) \frac{v^p}{u^{p-1}} \, dz + \int_{\partial\Omega} \beta(z) u^{p-1} \frac{v^p}{u^{p-1}} \, d\sigma \text{ (via the nonlinear Green identity, see [6, p. 211])} \\ &\leq \|Dv\|_p^p - \lambda \|v\|_p^p + \int_{\partial\Omega} \beta(z) v^p \, d\sigma \text{ (see (4.13))} \\ &= \tau(v) - \lambda \|v\|_p^p. \end{aligned}$$

Let $v = \hat{u}_1 \in D_+$. Then

$$0 \leq [\hat{\lambda}_1 - \lambda] < 0 \text{ (since } \lambda > \hat{\lambda}_m, m \geq 2 \text{ and } \|\hat{u}_1\|_p = 1),$$

a contradiction. This proves Claim 4.3.

The homotopy invariance property of critical groups (see Gasinski & Papageorgiou [10, Theorem 5.125, p. 836]) implies that for sufficiently small $r > 0$ we have

$$H_k((\hat{h}_\lambda^+)(0, \cdot)^\circ \cap B_r, (\hat{h}_\lambda^+)(0, \cdot)^\circ \cap B_r \setminus \{0\}) = H_k((\hat{h}_\lambda^+)(1, \cdot)^\circ \cap B_r, (\hat{h}_\lambda^+)(1, \cdot)^\circ \cap B_r \setminus \{0\}) \text{ for all } k \in \mathbf{N}_0. \quad (4.14)$$

On account of Claim 4.3, 0 is an ordinary level for $\hat{h}_\lambda^+(1, \cdot)$. Hence from Granas & Dugundji [12, p. 407], we have

$$H_k((\hat{h}_\lambda^+)(1, \cdot)^\circ \cap B_r, (\hat{h}_\lambda^+)(1, \cdot)^\circ \cap B_r \setminus \{0\}) = 0 \text{ for all } k \in \mathbf{N}_0. \quad (4.15)$$

From the definition of critical groups, we have

$$H_k((\hat{h}_\lambda^+)(0, \cdot)^\circ \cap B_r, (\hat{h}_\lambda^+)(0, \cdot)^\circ \cap B_r \setminus \{0\}) = C_k(\psi_\lambda^+, 0) \text{ for all } k \in \mathbf{N}_0. \quad (4.16)$$

Combining (4.14), (4.15), (4.16), we obtain

$$C_k(\psi_\lambda^+, 0) = 0 \text{ for all } k \in \mathbf{N}_0. \quad (4.17)$$

Since $\lambda \in (\hat{\lambda}_m, \hat{\lambda}_{m+1}) \setminus \hat{\sigma}(p)$, we have

$$\begin{aligned} K_{\psi_\lambda^+} &= \{0\}, \\ \Rightarrow C_k(\psi_\lambda^+, 0) &= C_k(\psi_\lambda^+, \infty) \text{ for all } k \in \mathbf{N}_0 \text{ (see [19, Proposition 6.61, p. 160]).} \end{aligned} \quad (4.18)$$

By (4.11), (4.17), (4.18), we can conclude that

$$C_k(\chi_\lambda^+, \infty) = 0 \text{ for all } k \in \mathbf{N}_0.$$

Similarly, we can show that

$$C_k(\chi_\lambda^-, \infty) = 0 \text{ for all } k \in \mathbf{N}_0.$$

The proof is now complete. □

Let $\hat{u} \in D_+$ and $\hat{v} \in -D_+$ be the second pair of constant sign solutions for problem (P_λ) ($\lambda > 0$ sufficiently large) produced in Proposition 3.4.

Proposition 4.4. *If hypotheses $H(\beta)$, $H(g)$, $H(f)$, H_0 hold and $\lambda > 0$ is large enough (see Proposition 3.4), then $C_k(\chi_\lambda^+, \hat{u}) = C_k(\chi_\lambda^-, \hat{v}) = \delta_{k,1}\mathbf{Z}$ for all $k \in \mathbf{N}_0$.*

Proof. We may assume that $K_{\chi_\lambda^+} = \{u_0, \hat{u}\}$. Otherwise we already have a fifth nontrivial solution for (P_λ) , which is also positive (see (3.46) and (3.25)).

Let $\hat{m}_\lambda^+ = \chi_\lambda^+(u_0)$ and let \tilde{m}_λ^+ be as in (3.44). We have $\hat{m}_\lambda^+ < \tilde{m}_\lambda^+$ and we choose $\eta, \vartheta \in \mathbf{R}$ such that

$$\eta < \hat{m}_\lambda^+ < \vartheta < \tilde{m}_\lambda^+. \tag{4.19}$$

For these levels, we consider the corresponding sublevel sets for χ_λ^+

$$(\chi_\lambda^+)^{\eta} \subseteq (\chi_\lambda^+)^{\vartheta} \subseteq W^{1,p}(\Omega).$$

For this triple we consider the corresponding long exact sequence of singular homological groups (see Motreanu, Motreanu & Papageorgiou [19, Proposition 6.14, p. 143]). We have

$$\dots \rightarrow H_k(W^{1,p}(\Omega), (\chi_\lambda^+)^{\eta}) \xrightarrow{i_*} H_k(W^{1,p}(\Omega), (\chi_\lambda^+)^{\vartheta}) \xrightarrow{\hat{\partial}_*} H_{k-1}((\chi_\lambda^+)^{\eta}, (\chi_\lambda^+)^{\vartheta}) \rightarrow \dots \tag{4.20}$$

with i_* being the homomorphism induced by the inclusion map $i : (W^{1,p}(\Omega), (\chi_\lambda^+)^{\eta}) \rightarrow (W^{1,p}(\Omega), (\chi_\lambda^+)^{\vartheta})$ and $\hat{\partial}_*$ is the composite boundary homomorphism.

From (4.19) we see that $\eta < \inf \chi_\lambda^+(K_{\chi_\lambda^+})$ and so

$$H_k(W^{1,p}(\Omega), (\chi_\lambda^+)^{\eta}) = C_k(\chi_\lambda^+, \infty) = 0 \quad \text{for all } k \in \mathbf{N}_0 \text{ (see Proposition 4.1).} \tag{4.21}$$

Also, from (4.19) and (3.46), we have

$$H_k(W^{1,p}(\Omega), (\chi_\lambda^+)^{\vartheta}) = C_k(\chi_\lambda^+, \hat{u}) \quad \text{for all } k \in \mathbf{N}_0, \tag{4.22}$$

$$H_{k-1}((\chi_\lambda^+)^{\eta}, (\chi_\lambda^+)^{\vartheta}) = C_{k-1}(\chi_\lambda^+, u_0) = \delta_{k-1,0}\mathbf{Z} = \delta_{k,1}\mathbf{Z} \quad \text{for all } k \in \mathbf{N}_0 \tag{4.23}$$

(see [19, Lemma 6.55, p. 175] and Claim 2 of Proposition 3.4).

Returning to (4.20) and using (4.21), (4.22), (4.23), we see that only the tail (that is, $k = 1$) of the long exact sequence is nontrivial. Moreover, by the rank theorem and the exactness of (4.20), we have

$$\begin{aligned} \text{rank } H_1(W^{1,p}(\Omega), (\chi_\lambda^+)^{\vartheta}) &= \text{rank ker } \hat{\partial}_* + \text{rank im } \hat{\partial}_* = \text{rank im } i_* + \text{rank im } \hat{\partial}_*, \\ &\Rightarrow \text{rank } C_1(\chi_\lambda^+, \hat{u}) \leq 1 \quad \text{(see (4.21), (4.22), (4.23)).} \end{aligned} \tag{4.24}$$

From the proof of Proposition 3.4 we know that $\hat{u} \in K_{\chi_\lambda^+}$ is of the mountain pass type. Therefore

$$C_1(\chi_\lambda^+, \hat{u}) \neq 0 \tag{4.25}$$

(see [19, Corollary 6.81, p. 168]).

It follows from (4.24) and (4.25) that

$$C_k(\chi_\lambda^+, \hat{u}) = \delta_{k,1}\mathbf{Z} \quad \text{for all } k \in \mathbf{N}_0.$$

Similarly, we can show that

$$C_k(\chi_\lambda^-, \hat{v}) = \delta_{k,1}\mathbf{Z} \quad \text{for all } k \in \mathbf{N}_0.$$

This completes the proof. □

Let $\varphi_\lambda : W^{1,p}(\Omega) \rightarrow \mathbf{R}$ be the energy functional for problem (P_λ) defined by

$$\varphi_\lambda(u) = \frac{1}{p} \tau(u) - \lambda \int_{\Omega} G(z, u) dz - \int_{\Omega} F(z, u) dz \quad \text{for all } u \in W^{1,p}(\Omega).$$

Evidently, $\varphi_\lambda \in C^1(W^{1,p}(\Omega), \mathbf{R})$.

Proposition 4.5. *If hypotheses $H(\beta)$, $H(g)$, $H(f)$ hold and $\lambda > 0$, then the functional φ_λ satisfies the C-condition.*

Proof. Let $\{u_n\}_{n \geq 1} \subseteq W^{1,p}(\Omega)$ be a sequence such that

$$|\varphi_\lambda(u_n)| \leq M_2 \text{ for some } M_2 > 0 \text{ and all } n \in \mathbf{N}, \quad (4.26)$$

$$(1 + \|u_n\|)\varphi'_\lambda(u_n) \rightarrow 0 \text{ in } W^{1,p}(\Omega)^* \text{ as } n \rightarrow \infty. \quad (4.27)$$

By (4.27) we have

$$\left| \langle A(u_n), h \rangle + \int_{\partial\Omega} \beta(z) |u_n|^{p-2} u_n h d\sigma - \lambda \int_{\Omega} g(z, u_n) h dz - \int_{\Omega} f(z, u_n) h dz \right| \leq \frac{\epsilon_n \|h\|}{1 + \|u_n\|} \quad (4.28)$$

for all $h \in W^{1,p}(\Omega)$ with $\epsilon_n \rightarrow 0^+$.

In (4.28) we choose $h = u_n \in W^{1,p}(\Omega)$. Then

$$\tau(u_n) - \int_{\Omega} [\lambda g(z, u_n) + f(z, u_n)] u_n dz \leq \epsilon_n \quad \text{for all } n \in \mathbf{N}. \quad (4.29)$$

Also, from (4.26) we have

$$-\tau(u_n) + \int_{\Omega} p[\lambda G(z, u_n) + F(z, u_n)] dz \leq pM_2 \quad \text{for all } n \in \mathbf{N}. \quad (4.30)$$

Adding (4.29) and (4.30), we obtain

$$\int_{\Omega} [p(\lambda G(z, u_n) + F(z, u_n)) - (\lambda g(z, u_n) + f(z, u_n)) u_n] dz \leq M_3 \text{ for some } M_3 > 0 \text{ and all } n \in \mathbf{N}. \quad (4.31)$$

We claim that $\{u_n\}_{n \geq 1} \subseteq W^{1,p}(\Omega)$ is bounded. Arguing by contradiction, suppose that $\|u_n\| \rightarrow \infty$. We set $y_n = \frac{u_n}{\|u_n\|}$, $n \in \mathbf{N}$. We have $\|y_n\| = 1$ for all $n \in \mathbf{N}$ and so we may assume that

$$y_n \xrightarrow{w} y \text{ in } W^{1,p}(\Omega) \quad \text{and} \quad y_n \rightarrow y \text{ in } L^p(\Omega) \text{ and } L^p(\partial\Omega). \quad (4.32)$$

From (4.28) we have

$$\left| \langle A(y_n), h \rangle + \int_{\partial\Omega} \beta(z) |y_n|^{p-2} y_n h d\sigma - \int_{\Omega} \frac{\lambda g(z, u_n)}{\|u_n\|^{p-1}} h dz - \int_{\Omega} \frac{f(z, u_n)}{\|u_n\|^{p-1}} h dz \right| \leq \frac{\epsilon_n \|h\|}{(1 + \|u_n\|) \|u_n\|^{p-1}} \quad (4.33)$$

for all $n \in \mathbf{N}$.

From (2.5) it is clear that

$$\left\{ \frac{g(\cdot, u_n(\cdot))}{\|u_n\|^{p-1}} \right\}_{n \in \mathbf{N}}, \left\{ \frac{f(\cdot, u_n(\cdot))}{\|u_n\|^{p-1}} \right\}_{n \in \mathbf{N}} \subseteq L^{p'}(\Omega) \text{ are bounded sequences.} \quad (4.34)$$

In (4.33) we choose $h = y_n - y \in W^{1,p}(\Omega)$ and pass to the limit as $n \rightarrow \infty$. Then using (4.32) and (4.34), we obtain

$$\lim_{n \rightarrow \infty} \langle A(y_n), y_n - y \rangle = 0, \quad \Rightarrow y_n \rightarrow y \text{ in } W^{1,p}(\Omega) \text{ (see Proposition 2.4) and so } \|y\| = 1. \quad (4.35)$$

From (4.35) we see that $y \neq 0$ and so if $D_+ = \{z \in \Omega : |y(z)| > 0\}$, then $|D_+|_N > 0$ with $|\cdot|_N$ denoting the Lebesgue measure on \mathbf{R}^N . We have

$$\begin{aligned} &|u_n(z)| \rightarrow +\infty \text{ for almost all } z \in D_+, \\ &\Rightarrow \liminf_{n \rightarrow \infty} \frac{pF(z, u_n(z)) - f(z, u_n(z))u_n(z)}{|u_n(z)|^\tau} > 0 \text{ for almost all } z \in D_+ \quad (\text{see hypothesis } H(f)(iii)) \\ &\Rightarrow \liminf_{n \rightarrow \infty} \frac{1}{\|u_n\|^\tau} \int_{D_+} [pF(z, u_n) - f(z, u_n)u_n] dz > 0 \text{ (by Fatou's lemma)} \\ &\Rightarrow \liminf_{n \rightarrow \infty} \frac{1}{\|u_n\|^\tau} \int_{\Omega} [pF(z, u_n) - f(z, u_n)u_n] dz > 0 \text{ (see } H(f)(ii)). \end{aligned} \tag{4.36}$$

Note that hypothesis $H(g)(ii)$ implies that given $\epsilon > 0$, we can find $c_{17} = c_{17}(\epsilon) > 0$ such that

$$g(z, x)x \leq \epsilon|x|^\tau + c_{17} \text{ for almost all } z \in \Omega \text{ and all } x \in \mathbf{R} \text{ (see hypothesis } H(g)(i)). \tag{4.37}$$

Since $G(z, x) \geq 0$ for almost all $z \in \Omega$, all $x \in \mathbf{R}$ (by the sign condition in $H(g)(i)$), we obtain

$$pG(z, x) - g(z, x)x \geq -\epsilon|x|^\tau - c_{17} \text{ for almost all } z \in \Omega \text{ and all } x \in \mathbf{R} \text{ (see (4.37)).} \tag{4.38}$$

Hence

$$\begin{aligned} &\int_{\Omega} [p(\lambda G(z, u_n) + F(z, u_n)) - (\lambda g(z, u_n) + f(z, u_n))u_n] dz \\ &\geq \int_{\Omega} [-\lambda\epsilon|u_n|^\tau + (pF(z, u_n) - f(z, u_n)u_n)] dz \text{ (see (4.38))} \\ &\Rightarrow \frac{1}{\|u_n\|^\tau} \int_{\Omega} [p(\lambda G(z, u_n) + F(z, u_n)) - (\lambda g(z, u_n) + f(z, u_n))u_n] dz \\ &\geq \int_{\Omega} \frac{-\lambda\epsilon}{\|u_n\|^{p-\tau}} |y_n|^\tau dz + \frac{1}{\|u_n\|^p} \int_{\Omega} [pF(z, u_n) - f(z, u_n)u_n] dz. \end{aligned}$$

Using (4.36), we obtain

$$\liminf_{n \rightarrow \infty} \frac{1}{\|u_n\|^\tau} \int_{\Omega} [p(\lambda G(z, u_n) + F(z, u_n)) - (\lambda g(z, u_n) + f(z, u_n))u_n] dz > 0. \tag{4.39}$$

On the other hand, relation (4.31) yields

$$\limsup_{n \rightarrow \infty} \frac{1}{\|u_n\|^\tau} \int_{\Omega} [p(\lambda G(z, u_n) + F(z, u_n)) - (\lambda g(z, u_n) + f(z, u_n))u_n] dz \leq 0. \tag{4.40}$$

Comparing (4.39) and (4.40), we arrive at a contradiction.

This proves that

$$\{u_n\}_{n \geq 1} \subseteq W^{1,p}(\Omega) \text{ is bounded.}$$

So, we may assume that

$$u_n \overset{w}{\rightharpoonup} u \text{ in } W^{1,p}(\Omega) \quad \text{and} \quad u_n \rightarrow u \text{ in } L^p(\Omega) \text{ and } L^p(\partial\Omega). \tag{4.41}$$

In (4.28) we choose $h = u_n - u \in W^{1,p}(\Omega)$, pass to the limit as $n \rightarrow \infty$, and use (4.41). Then

$$\begin{aligned} &\lim_{n \rightarrow \infty} \langle A(u_n), u_n - u \rangle = 0, \\ &\Rightarrow u_n \rightarrow u \text{ in } W^{1,p}(\Omega) \text{ (see Proposition 2.4),} \\ &\Rightarrow \varphi_\lambda \text{ satisfies the } C\text{-condition.} \end{aligned}$$

The proof is now complete. □

Then using Proposition 8 of Papageorgiou, Rădulescu & Repovš [25], (see also [4]), we obtain the following property.

Proposition 4.6. *If hypotheses $H(\beta)$, $H(g)$, $H(f)$ hold and $\lambda > 0$, then $C_m(\varphi_\lambda, \infty) \neq 0$.*

We assume that K_{φ_λ} ($\lambda > 0$ sufficiently large, as in Proposition 3.4) is finite. Otherwise we already have an infinity of solutions which are in $C^1(\overline{\Omega})$ (nonlinear regularity theory).

Proposition 4.7. *If hypotheses $H(\beta)$, $H(g)$, $H(f)$, H_0 hold and $\lambda > 0$ is sufficiently large (see Proposition 3.4), then*

$$C_k(\varphi_\lambda, \hat{u}) = C_k(\varphi_\lambda, \hat{v}) = \delta_{k,1} \mathbf{Z} \quad \text{for all } k \in \mathbf{N}_0,$$

$$C_k(\varphi_\lambda, 0) = C_k(\varphi_\lambda, u_0) = C_k(\varphi_\lambda, v_0) = \delta_{k,0} \mathbf{Z} \quad \text{for all } k \in \mathbf{N}_0.$$

Proof. Consider the homotopy $\tilde{h}_\lambda^+(\cdot, \cdot)$ defined by

$$\tilde{h}_\lambda^+(t, u) = (1 - t)\varphi_\lambda(u) + t\chi_\lambda^+(u) \quad \text{for all } (t, u) \in [0, 1] \times W^{1,p}(\Omega).$$

Suppose that we can find $t_n \rightarrow t$ and $u_n \rightarrow \hat{u}$ in $W^{1,p}(\Omega)$ such that

$$(\tilde{h}_\lambda^+)'_n(t_n, u_n) = 0 \quad \text{for all } n \in \mathbf{N}.$$

We have

$$\begin{aligned} & \langle A(u_n), h \rangle + \int_{\partial\Omega} \beta(z)|u_n|^{p-2}u_n h \, d\sigma \\ &= (1 - t_n) \int_{\Omega} (\lambda g(z, u_n) + f(z, u_n))h \, dz + t_n \int_{\Omega} i_\lambda^+(z, u_n)h \, dz \quad \text{for all } h \in W^{1,p}(\Omega), n \in \mathbf{N}, \\ &\Rightarrow -\Delta_p u_n(z) = (1 - t_n)(\lambda g(z, u_n(z)) + f(z, u_n(z))) + t_n i_\lambda^+(z, u_n(z)) \quad \text{for almost all } z \in \Omega, \\ &\frac{\partial u}{\partial n_p} + \beta(z)|u_n|^{p-2}u_n = 0 \text{ on } \partial\Omega \text{ (see Papageorgiou \& R\u0105dulescu [21])}. \end{aligned} \tag{4.42}$$

From (4.42) and Proposition 7 of Papageorgiou & Rădulescu [23], we have

$$\|u_n\|_\infty \leq M_4 \text{ for some } M_4 > 0 \text{ and all } n \in \mathbf{N}.$$

Then invoking Theorem 2 of Lieberman [17], we can find $\alpha \in (0, 1)$ and $M_5 > 0$ such that

$$u_n \in C^{1,\alpha}(\overline{\Omega}) \text{ and } \|u_n\|_{C^{1,\alpha}(\overline{\Omega})} \leq M_5 \text{ for some } M_5 > 0 \text{ and all } n \in \mathbf{N}. \tag{4.43}$$

By (4.43), the compact embedding of $C^{1,\alpha}(\overline{\Omega})$ into $C^1(\overline{\Omega})$ and the fact that $u_n \rightarrow \hat{u}$ in $W^{1,p}(\Omega)$, we infer that

$$\begin{aligned} & u_n \rightarrow \hat{u} \text{ in } C^1(\overline{\Omega}), \\ & \Rightarrow u_n - u_0 \in \text{int } C_+ \quad \text{for all } n \geq n_0 \text{ (see Proposition 3.4),} \\ & \Rightarrow \{u_n\}_{n \geq n_0} \subseteq K_{\varphi_\lambda} \text{ (see (3.25)),} \end{aligned}$$

a contradiction to our hypothesis that K_{φ_λ} is finite.

Therefore by the homotopy invariance property of critical groups (see [10, p. 836]), we have

$$C_k(\varphi_\lambda, \hat{u}) = C_k(\chi_\lambda^+, \hat{u}) \quad \text{for all } k \in \mathbf{N}_0, \quad \Rightarrow C_k(\varphi_\lambda, \hat{u}) = \delta_{k,1} \mathbf{Z} \quad \text{for all } k \in \mathbf{N}_0.$$

Similarly, using this time χ_λ^- , we show that

$$C_k(\varphi_\lambda, \hat{v}) = \delta_{k,1} \mathbf{Z} \quad \text{for all } k \in \mathbf{N}_0.$$

Recall that $u_0 \in D_+$ and $v_0 \in -D_+$ are local minimizers of the functionals $\chi_\lambda^+(\cdot)$ and $\chi_\lambda^-(\cdot)$, respectively (see Claim 4.3 in the proof of Proposition 3.4). Hence we have

$$C_k(\chi_\lambda^+, u_0) = C_k(\chi_\lambda^-, v_0) = \delta_{k,0} \mathbf{Z} \quad \text{for all } k \in \mathbf{N}_0. \quad (4.44)$$

A homotopy invariance argument as above, shows that

$$\begin{aligned} C_k(\varphi_\lambda, u_0) &= C_k(\chi_\lambda^+, u_0) \text{ and } C_k(\varphi_\lambda, v_0) = C_k(\chi_\lambda^-, v_0) \quad \text{for all } k \in \mathbf{N}_0, \\ &\Rightarrow C_k(\varphi_\lambda, u_0) = C_k(\varphi_\lambda, v_0) = \delta_{k,0} \mathbf{Z} \quad \text{for all } k \in \mathbf{N}_0 \text{ (see (4.44)).} \end{aligned}$$

Finally, hypotheses $H(g)(ii)$ and $H(f)(iii)$ imply that

$$u = 0 \text{ is a local minimizer of } \varphi_\lambda$$

(see also the proof of Proposition 3.1). It follows that

$$C_k(\varphi_\lambda, 0) = \delta_{k,0} \mathbf{Z} \quad \text{for all } k \in \mathbf{N}_0.$$

The proof is now complete. □

Proposition 4.8. *If hypotheses $H(\beta)$, $H(g)$, $H(f)$, H_0 hold and $\lambda > 0$ is big (see Proposition 3.4), then problem (P_λ) has a fifth nontrivial solution*

$$y_0 \in C^1(\overline{\Omega}).$$

Proof. According to Proposition 4.6, we have

$$C_m(\varphi_\lambda, \infty) \neq 0.$$

So, there exists $y_0 \in K_{\varphi_\lambda}$ such that

$$C_m(\varphi_\lambda, y_0) \neq 0. \quad (4.45)$$

Since $m \geq 2$, by Proposition 4.7 and (4.45), we infer that

$$y_0 \notin \{0, u_0, v_0, \hat{u}, \hat{v}\}.$$

Therefore y_0 is a fifth nontrivial solution of (P_λ) (for sufficiently large $\lambda > 0$) and the nonlinear regularity theory implies that $y_0 \in C^1(\overline{\Omega})$. □

Finally, we can state the following multiplicity theorem for problem (P_λ) .

Theorem 4.9. *If hypotheses $H(\beta)$, $H(g)$, $H(f)$, H_0 hold, then for all sufficiently large $\lambda > 0$ problem (P_λ) has at least five nontrivial solutions*

$$u_0, \hat{u} \in D_+ \text{ with } \hat{u} - u_0 \in \text{int } \hat{C}_+, \quad v_0, \hat{v} \in -D_+ \text{ with } v_0 - \hat{v} \in \text{int } \hat{C}_+ \quad \text{and} \quad y_0 \in C^1(\overline{\Omega}).$$

Question. Is it possible, in the framework of the present paper, to generate nodal solutions for (P_λ) ?

ACKNOWLEDGEMENTS

This research was supported by the Slovenian Research Agency grants P1-0292, J1-8131, J1-7025, N1-0064, and N1-0083.

ORCID

Vicențiu D. Rădulescu  <https://orcid.org/0000-0003-4615-5537>

Dušan D. Repovš  <https://orcid.org/0000-0002-6643-1271>

REFERENCES

- [1] S. Aizicovici, N. S. Papageorgiou, and V. Staicu, *Degree theory for operators of monotone type and nonlinear elliptic equations with inequality constraints*, Mem. Amer. Math. Soc. **196** (2008), no. 915.
- [2] W. Allegretto and Y. X. Huang, *A Picone's identity for the p -Laplacian and applications*, Nonlinear Anal. **32** (1998), 819–830.
- [3] K. C. Chang, *Methods in nonlinear analysis*, Springer-Verlag, Berlin, 2005.
- [4] S. Cingolani and M. Degiovanni, *Nontrivial solutions for p -Laplacian equations with right-hand side having p -linear growth at infinity*, Comm. Partial Differential Equations **30** (2005), 1191–1203.
- [5] E. Fadell and P. Rabinowitz, *Generalized cohomological index theories for Lie group actions with applications to bifurcation questions for Hamiltonian systems*, Invent. Math. **45** (1978), 139–174.
- [6] L. Gasinski and N. S. Papageorgiou, *Nonlinear analysis*, Chapman & Hall/CRC, Boca Raton, FL, 2006.
- [7] L. Gasinski and N. S. Papageorgiou, *Existence of three nontrivial smooth solutions for nonlinear resonant Neumann problems driven by the p -Laplacian*, Z. Anal. Anwend. **29** (2010), 413–428.
- [8] L. Gasinski and N. S. Papageorgiou, *Anisotropic nonlinear Neumann problems*, Calc. Var. Partial Differential Equations **42** (2011), 323–354.
- [9] L. Gasinski and N. S. Papageorgiou, *Multiplicity of positive solutions for eigenvalue problems of $(p, 2)$ -equations*, Bound. Value Probl. **2012**, no. 152, 17 pp.
- [10] L. Gasinski and N. S. Papageorgiou, *Exercises in analysis. part 2: nonlinear analysis*, Springer, Cham, 2016.
- [11] L. Gasinski and N. S. Papageorgiou, *Positive solutions for the Robin p -Laplacian problem with competing nonlinearities*, Adv. Calc. Var. **12** (2019), 31–56.
- [12] A. Granas and J. Dugundji, *Fixed point theory*, Springer-Verlag, New York, 2003.
- [13] Y. Guo and J. Liu, *Solutions of p -sublinear p -Laplacian equation via Morse theory*, J. Lond. Math. Soc. (2) **72** (2005), 623–644.
- [14] S. Hu and N. S. Papageorgiou, *Handbook of multivalued analysis. Volume I: Theory*, Kluwer Academic Publishers, Dordrecht, 1997.
- [15] Q. Jiu and J. Su, *Existence and multiplicity results for Dirichlet problems with p -Laplacian*, J. Math. Anal. Appl. **281** (2003), 587–601.
- [16] Z. Liang and J. Su, *Multiple solutions for semilinear elliptic boundary value problems with double resonance*, J. Math. Anal. Appl. **354** (2009), 147–158.
- [17] G. Lieberman, *Boundary regularity for solutions of degenerate elliptic equations*, Nonlinear Anal. **12** (1988), 1203–1219.
- [18] S. Liu, *Multiple solutions for coercive p -Laplacian equations*, J. Math. Anal. Appl. **316** (2006), 229–236.
- [19] D. Motreanu, V. Motreanu, and N. S. Papageorgiou, *Topological and variational methods with applications to nonlinear boundary value problems*, Springer, New York, 2014.
- [20] D. Mugnai and N. S. Papageorgiou, *Resonant nonlinear Neumann problems with indefinite weight*, Annali Sc. Norm. Pisa, Cl. Sci. (5) (2012), 729–788.
- [21] N. S. Papageorgiou and V. D. Rădulescu, *Multiple solutions with precise sign for parametric Robin problems*, J. Differential Equations **256** (2014), 2449–2479.
- [22] N. S. Papageorgiou and V. D. Rădulescu, *Coercive and noncoercive nonlinear Neumann problems with indefinite potential*, Forum Math. **28** (2016), 545–571.
- [23] N. S. Papageorgiou and V. D. Rădulescu, *Nonlinear nonhomogeneous Robin problems with superlinear reaction*, Adv. Nonlinear Stud. **16** (2016), 737–764.
- [24] N. S. Papageorgiou, V. D. Rădulescu, and D. D. Repovš, *Positive solutions for perturbations of the Robin eigenvalue problem plus an indefinite potential*, Discrete Cont. Dynam. Syst. **37** (2017), 2589–2618.
- [25] N. S. Papageorgiou, V. D. Rădulescu, and D. D. Repovš, *Multiple solutions for resonant problems of the Robin p -Laplacian plus an indefinite potential*, Calc. Var. Partial Differential Equations **56** (2017), no. 3, Art. 63, 23 pp.
- [26] N. S. Papageorgiou, V. D. Rădulescu, and D. D. Repovš, *Resonant Robin problems driven by the p -Laplacian plus an indefinite potential*, Ann. Acad. Sci. Fenn. Math. **43** (2018), 483–508.
- [27] N. S. Papageorgiou, V. D. Rădulescu, and D. D. Repovš, *Nonlinear analysis – theory and methods*, Springer Monogr. Math., Springer, Cham, Switzerland, 2019.

How to cite this article: Papageorgiou NS, Rădulescu VD, Repovš DD. Parametric nonlinear resonant Robin problems. *Mathematische Nachrichten*. 2019;292:2456–2480. <https://doi.org/10.1002/mana.201800505>