



Anisotropic equations with indefinite potential and competing nonlinearities[☆]



Nikolaos S. Papageorgiou^{a,e}, Vicențiu D. Rădulescu^{b,c,e,*}, Dušan D. Repovš^d

^a National Technical University, Department of Mathematics, Zografou Campus, Athens 15780, Greece

^b Faculty of Applied Mathematics, AGH University of Science and Technology, al. Mickiewicza 30, 30-059 Kraków, Poland

^c Department of Mathematics, University of Craiova, 200585 Craiova, Romania

^d Faculty of Education and Faculty of Mathematics and Physics, University of Ljubljana, 1000 Ljubljana, Slovenia

^e Institute of Mathematics, Physics and Mechanics, 1000 Ljubljana, Slovenia

ARTICLE INFO

Article history:

Received 2 January 2020

Accepted 10 March 2020

Communicated by Enrico Valdinoci

MSC:

35J10

35J70

Keywords:

Variable exponent spaces

Regularity theory

Maximum principle

Concave and convex nonlinearities

Positive solutions

Comparison principles

ABSTRACT

We consider a nonlinear Dirichlet problem driven by a variable exponent p -Laplacian plus an indefinite potential term. The reaction has the competing effects of a parametric concave (sublinear) term and a convex (superlinear) perturbation (the anisotropic concave–convex problem). We prove a bifurcation-type theorem describing the changes in the set of positive solutions as the positive parameter λ varies. Also, we prove the existence of minimal positive solutions.

© 2020 Elsevier Ltd. All rights reserved.

1. Introduction

Let $\Omega \subseteq \mathbb{R}^N$ be a bounded domain with a C^2 -boundary $\partial\Omega$. In this paper we study the following anisotropic boundary value problem

$$\begin{cases} -\Delta_{p(z)}u(z) + \xi(z)u(z)^{p(z)-1} = \lambda u(z)^{q(z)-1} + f(z, u(z)) & \text{in } \Omega, \\ u|_{\partial\Omega} = 0, \lambda > 0, u > 0. \end{cases} \quad (P_\lambda)$$

[☆] This paper is dedicated with esteem to Professor Shair Ahmad on the occasion of his 85th anniversary.

* Corresponding author at: Faculty of Applied Mathematics, AGH University of Science and Technology, al. Mickiewicza 30, 30-059 Kraków, Poland.

E-mail addresses: npapg@math.ntua.gr (N.S. Papageorgiou), radulescu@inf.ucv.ro (V.D. Rădulescu), dusan.repovs@guest.arnes.si (D.D. Repovš).

In this problem, $\Delta_{p(z)}$ denotes the $p(z)$ -Laplacian defined by

$$\Delta_{p(z)}u = \operatorname{div}(|Du|^{p(z)-2}Du(z)) \text{ for all } u \in W_0^{1,p(z)}(\Omega).$$

Concerning the exponents $p, q : \overline{\Omega} \rightarrow \mathbb{R}$, we assume that both are functions belonging to $C^{0,1}(\overline{\Omega})$ and satisfying

$$1 < q_- \leq q(z) \leq q_+ < p_- \leq p(z) \leq p_+ \text{ for all } z \in \overline{\Omega}.$$

The potential function $\xi \in L^\infty(\Omega)$ is sign-changing. Therefore the differential operator of (P_λ) (left-hand side) is not coercive. In the reaction (the right-hand side of (P_λ)), we have a parametric term with $\lambda > 0$ being the parameter and a perturbation $f(z, x)$ which is jointly measurable and of class C^1 in the x -variable. We assume that $f(z, \cdot)$ exhibits $(p_+ - 1)$ -superlinear growth near $+\infty$ without satisfying the usual in such cases Ambrosetti–Rabinowitz condition (*AR*-condition for short). So, in the reaction of problem (P_λ) we have the competing effects of a sublinear (concave) term and of a superlinear (convex) term. We are looking for positive solutions and our aim is to obtain a precise description of the changes in the set of positive solutions as the parameter $\lambda > 0$ varies (a bifurcation-type result).

The study of such parametric concave–convex problems started with the seminal paper of Ambrosetti, Brezis and Cerami [1], where $p(z) = 2$ for all $z \in \overline{\Omega}$ (the semilinear isotropic problem). It was extended to equations driven by the p -Laplacian and with the reaction being $\lambda x^{q-1} + x^{r-1}$ for all $x \geq 0$ with $1 < q < p < r < p^*$ by Garcia Azorero, Manfredi and Peral Alonso [10], and Guo and Zhang [13]. Recall that

$$p^* = \begin{cases} \frac{Np}{N-p}, & \text{if } p \leq N \\ +\infty, & \text{if } N < p. \end{cases}$$

Further extensions can be found in the works of Marano and Papageorgiou [16] and Papageorgiou and Rădulescu [17]. All the aforementioned works deal with isotropic equations. To the best of our knowledge, no such results exist for anisotropic equations.

Additional parametric boundary value problems driven by operators with variable exponents and applications, can be found in the book of Rădulescu and Repovš [23]. We also refer to the recent papers [4,5,19,21,22,27], all dealing with isotropic or anisotropic nonlinear problems with Dirichlet boundary condition.

2. Mathematical background, auxiliary results and hypotheses

In this section we briefly review some basic facts about variable exponent spaces and we prove two anisotropic strong comparison theorems which we will need in our analysis of problem (P_λ) .

A comprehensive presentation of variable exponent Lebesgue and Sobolev spaces can be found in the book of Diening, Harjulehto, Hästö and Ruzička [6].

So, let $L_1^\infty(\Omega) = \{p \in L^\infty(\Omega) : \operatorname{ess\,inf}_\Omega p \geq 1\}$. For $p \in L_1^\infty(\Omega)$, we set

$$p_- = \operatorname{ess\,inf}_\Omega p \quad \text{and} \quad p_+ = \operatorname{ess\,sup}_\Omega p.$$

Also let $M(\Omega) = \{u : \Omega \rightarrow \mathbb{R} : u(\cdot) \text{ is measurable}\}$. As usual, we identify two functions which differ on a set of zero measure.

Given $p \in L_1^\infty(\Omega)$, we define the following variable exponent Lebesgue space

$$L^{p(z)}(\Omega) = \left\{ u \in M(\Omega) : \int_\Omega |u|^{p(z)} dz < +\infty \right\}.$$

We equip $L^{p(z)}(\Omega)$ with the following norm (known as the Luxemburg norm)

$$\|u\|_{p(z)} = \inf \left\{ \lambda > 0 : \int_\Omega \left(\frac{|u|}{\lambda} \right)^{p(z)} dz \leq 1 \right\}.$$

Having defined variable exponent Lebesgue spaces, we can introduce variable exponent Sobolev spaces by

$$W^{1,p(z)}(\Omega) = \{u \in L^{p(z)}(\Omega) : |Du| \in L^{p(z)}(\Omega)\}.$$

We equip this space with the following norm

$$\|u\|_{1,p(z)} = \|u\|_{p(z)} + \|Du\|_{p(z)}.$$

An equivalent norm of $W^{1,p(z)}(\Omega)$ is given by

$$\|u\|'_{1,p(z)} = \inf \left\{ \lambda > 0 : \int_{\Omega} \left(\left(\frac{|Du|}{\lambda} \right)^{p(z)} + \left(\frac{|u|}{\lambda} \right)^{p(z)} \right) dz \leq 1 \right\}.$$

We define $W_0^{1,p(z)}(\Omega)$ as the closure in the $\|\cdot\|_{1,p(z)}$ of all compactly supported $W^{1,p(z)}(\Omega)$ -functions.

When $p \in L_1^\infty(\Omega)$ and $p_- > 1$, then the spaces $L^{p(z)}(\Omega)$, $W^{1,p(z)}(\Omega)$ and $W_0^{1,p(z)}(\Omega)$ are all separable, reflexive and uniformly convex.

We set

$$p^*(z) = \begin{cases} \frac{Np(z)}{N-p(z)}, & \text{if } p(z) < N \\ +\infty, & \text{if } p(z) \geq N. \end{cases}$$

If $p, q \in C(\bar{\Omega})$, $p_+ < N$ and $1 \leq q(z) \leq p^*(z)$ (resp. $1 \leq q(z) < p^*(z)$) for all $z \in \bar{\Omega}$, then $W^{1,p(z)}(\Omega)$ and $W_0^{1,p(z)}(\Omega)$ are embedded continuously (resp. compactly) into $L^{q(z)}(\Omega)$.

If $p, p' \in L_1^\infty(\Omega)$ and $\frac{1}{p} + \frac{1}{p'} = 1$, then $L^{p(z)}(\Omega)^* = L^{p'(z)}(\Omega)$ and we have the following Hölder type inequality

$$\int_{\Omega} |uv| dz \leq \left(\frac{1}{p_-} + \frac{1}{p'_-} \right) \|u\|_{p(z)} \|v\|_{p'(z)} \text{ for all } u \in L^{p(z)}(\Omega), v \in L^{p'(z)}(\Omega).$$

We say that $p \in C(\bar{\Omega})$ is logarithmic Hölder continuous (denoted by $p \in C^{0, \frac{1}{|\ln|\cdot|}}$) if it satisfies

$$|p(z) - p(z')| \leq \frac{c}{|\ln|z - z'||} \text{ for some } c > 0, \text{ all } z, z' \in \Omega, |z - z'| \leq \frac{1}{2}.$$

Note that $C^{0,1}(\bar{\Omega}) \hookrightarrow C^{0, \frac{1}{|\ln|\cdot|}}(\bar{\Omega})$. Also, when $p \in C^{0, \frac{1}{|\ln|\cdot|}}(\bar{\Omega})$, then

$$W_0^{1,p(z)}(\Omega) = \overline{C_c^\infty(\bar{\Omega})}^{\|\cdot\|_{1,p(z)}}.$$

Moreover, in this case the Poincaré inequality holds and we have

$$\|u\|_{p(z)} \leq \hat{C} \|Du\|_{p(z)} \text{ for all } u \in W_0^{1,p(z)}(\Omega),$$

where $\hat{C} > 0$ depends only on $(p, N, |\Omega|_N, \text{diam } \Omega)$, with $|\cdot|_N$ denoting the Lebesgue measure on \mathbb{R}^N . So, when $p \in C^{0, \frac{1}{|\ln|\cdot|}}(\bar{\Omega})$, then on the Sobolev space $W_0^{1,p(z)}(\Omega)$, we can use the equivalent norm

$$\|u\| = \|Du\|_{p(z)} \text{ for all } u \in W_0^{1,p(z)}(\Omega).$$

We introduce the following modular functions

$$\begin{aligned} \rho(u) &= \int_{\Omega} |u|^{p(z)} dz \text{ for all } u \in L^{p(z)}(\Omega), \\ \hat{\rho}(Du) &= \int_{\Omega} |Du|^{p(z)} dz \text{ for all } u \in W_0^{1,p(z)}(\Omega). \end{aligned}$$

We have the following property.

Proposition 2.1. (a) For every $u \in L^{p(z)}(\Omega)$, $u \neq 0$, we have

$$\|u\|_{p(z)} = \lambda \Leftrightarrow \rho\left(\frac{u}{\lambda}\right) = 1;$$

- (b) $\|u\|_{p(z)} < 1$ (resp. $= 1, > 1$) $\Leftrightarrow \rho(u) < 1$ (resp. $= 1, > 1$);
 (c) $\|u\|_{p(z)} < 1 \Rightarrow \|u\|_{p(z)}^{p_+} \leq \rho(u) \leq \|u\|_{p(z)}^{p_-}$ and $\|u\|_{p(z)} > 1 \Rightarrow \|u\|_{p(z)}^{p_-} \leq \rho(u) \leq \|u\|_{p(z)}^{p_+}$;
 (d) $\|u_n\|_{p(z)} \rightarrow 0 \Leftrightarrow \rho(u_n) \rightarrow 0$;
 (e) $\|u_n\|_{p(z)} \rightarrow +\infty \Leftrightarrow \rho(u_n) \rightarrow +\infty$.

Similarly, we have the following implications, when $p \in C^{0, \frac{1}{|\ln t|}}(\overline{\Omega})$.

Proposition 2.2. (a) For every $u \in W_0^{1,p(z)}(\Omega)$, $u \neq 0$, we have

$$\|u\| = \lambda \Leftrightarrow \hat{\rho}\left(\frac{Du}{\lambda}\right) = 1;$$

- (b) $\|u\| < 1$ (resp. $= 1, > 1$) $\Leftrightarrow \hat{\rho}(Du) < 1$ (resp. $= 1, > 1$);
 (c) $\|u\| < 1 \Rightarrow \|u\|^{p_+} \leq \hat{\rho}(Du) \leq \|u\|^{p_-}$ and $\|u\| > 1 \Rightarrow \|u\|^{p_-} \leq \hat{\rho}(Du) \leq \|u\|^{p_+}$;
 (d) $\|u_n\| \rightarrow 0 \Leftrightarrow \hat{\rho}(Du_n) \rightarrow 0$; (e) $\|u_n\| \rightarrow +\infty \Leftrightarrow \hat{\rho}(Du_n) \rightarrow +\infty$.

Let $p \in C^{0, \frac{1}{|\ln t|}}(\overline{\Omega})$. Then

$$W_0^{1,p(z)}(\Omega)^* = W^{-1,p'(z)}(\Omega) \quad \left(\frac{1}{p(z)} + \frac{1}{p'(z)} = 1\right).$$

Consider the operator $A : W_0^{1,p(z)}(\Omega) \rightarrow W^{-1,p'(z)}(\Omega) = W_0^{1,p(z)}(\Omega)^*$ defined by

$$\langle A(u), h \rangle = \int_{\Omega} |Du|^{p(z)-2} (Du, Dh)_{\mathbb{R}^N} dz \text{ for all } u, h \in W_0^{1,p(z)}(\Omega).$$

This operator has the following properties (see Gasinski and Papageorgiou [11]).

Proposition 2.3. The map $A : W_0^{1,p(z)}(\Omega) \rightarrow W^{-1,p'(z)}(\Omega)$ defined above is bounded (that is, maps bounded sets to bounded sets), continuous, strictly monotone (hence maximal monotone, too) and of type $(S)_+$, that is $u_n \xrightarrow{w} u$ in $W_0^{1,p(z)}(\Omega)$ and $\limsup_{n \rightarrow \infty} \langle A(u_n), (u_n - u) \rangle \leq 0 \Rightarrow u_n \rightarrow u$ in $W_0^{1,p(z)}(\Omega)$.

Next, we prove two strong comparison theorems, which will be used in the analysis of problem (P_λ) . The first one extends Proposition 2.6 of Arcoya and Ruiz [3] to the abstract setting of anisotropic problems.

We will use the following notation. Given $h, g \in L^\infty(\Omega)$, we write that $h \prec g$ if and only if for every $K \subseteq \Omega$ compact, we can find $c_K > 0$ such that $0 < c_K \leq g(z) - h(z)$ for a.a. $z \in K$. Evidently, if $h, g \in C(\Omega)$ and $h(z) < g(z)$ for all $z \in \Omega$, then $h \prec g$. Also, by C_+ we denote the positive cone of $C_0^1(\overline{\Omega}) = \{u \in C^1(\overline{\Omega}) : u|_{\partial\Omega} = 0\}$, that is, $C_+ = \{u \in C_0^1(\overline{\Omega}) : u(z) \geq 0 \text{ for all } z \in \overline{\Omega}\}$. This cone has a nonempty interior given by

$$\text{int } C_+ = \left\{ u \in C_+ : u(z) > 0 \text{ for all } z \in \Omega, \frac{\partial u}{\partial n} \Big|_{\partial\Omega} < 0 \right\},$$

with $n(\cdot)$ being the outward unit normal on $\partial\Omega$.

Proposition 2.4. *If $p \in C^{0,1}(\overline{\Omega})$, $1 < p(z)$ for all $z \in \overline{\Omega}$, $\hat{\xi}, h, g \in L^\infty(\Omega)$, $\hat{\xi}(z) \geq 0$ for a.a. $z \in \Omega$, $h \prec g$, $u \in W^{1,p(z)}(\Omega)$, $u \neq 0$, $v \in \text{int } C_+$ and*

$$\begin{aligned} & -\Delta_{p(z)}u + \hat{\xi}(z)|u|^{p(z)-2}u = h(z) \text{ in } \Omega, \quad u \Big|_{\partial\Omega} \leq 0, \\ & -\Delta_{p(z)}v + \hat{\xi}(z)v^{p(z)-1} = g(z) \text{ in } \Omega, \quad \frac{\partial v}{\partial n} \Big|_{\partial\Omega} < 0, \end{aligned}$$

then $v - u \in \text{int } C_+$.

Proof. From Theorem 4.1 of Fan and Zhao [8] (see also Proposition 3.1 of Gasinski and Papageorgiou [11]), we have that $u \in L^\infty(\Omega)$. Then invoking Theorem 1.3 of Fan [7], we infer that $u \in C^1(\overline{\Omega})$. Also, exploiting the monotonicity of $A(\cdot)$ (see Proposition 2.3), we see that $u \leq v$. We introduce the following two sets.

$$E = \{z \in \Omega : u(z) = v(z)\} \text{ and } \hat{E} = \{z \in \Omega : Du(z) = Dv(z) = 0\}.$$

Claim. $E \subseteq \hat{E}$.

Let $y = u - v$. We have $y \leq 0$. Consider $z \in E$. Then $y(z) = \max_\Omega y = 0$. So, we have $Dy(z) = 0$, hence $Du(z) = Dv(z)$. Arguing by contradiction, suppose that $z \notin \hat{E}$. Then $Dv(z) \neq 0$ and so we can find an open ball $B \subseteq \Omega$ centered at z such that

$$|Du(x)| > 0, \quad |Dv(x)| > 0, \quad (Du(x), Dv(x))_{\mathbb{R}^N} > 0 \text{ for all } x \in B.$$

Consider the $N \times N$ matrix $A(x) = (a_{ij}(x))_{i,j=1}^N$ with entries $a_{ij}(x)$ defined by

$$a_{ij}(x) = \int_0^1 [(1-t)Du(x) + tDv(x)] \left[\delta_{ij} + (p(x) - 2) \frac{D_i((1-t)u + tv)D_j((1-t)u + tv)}{|(1-t)Du + tDv|^2} \right] dt$$

We have $a_{ij} \in C^{0,\alpha}(\overline{B})$ for some $\alpha \in (0, 1)$ (see Fan [7]) and

$$-\text{div}(A(x)Dy(x)) = h(x) - g(x) - \hat{\xi}(x) \left[|u(x)|^{p(x)-2}u(x) - v(x)^{p(x)-1} \right] \text{ in } B \tag{1}$$

(see also Guedda and Véron [12]). By choosing the ball $B \subseteq \Omega$ even smaller if necessary, we obtain that in (1) the linear differential operator is strictly elliptic, while the right-hand side is strictly negative. Invoking Theorem 4 of Vázquez [25], we have

$$\begin{aligned} & y(x) < 0 \text{ for all } x \in B, \\ & \Rightarrow y(z) < 0, \text{ a contradiction.} \end{aligned}$$

Therefore $z \in \hat{E}$ and this proves the Claim.

Recall that $v \in \text{int } C_+$. Therefore \hat{E} is compact and so we can find $U \subseteq \Omega$ such that

$$\hat{E} \subseteq U \subseteq \overline{U} \subseteq \Omega. \tag{2}$$

For $\varepsilon > 0$ small enough, we have

$$u(z) + \varepsilon < v(z) \text{ for all } z \in \partial U \text{ (see (2))}, \tag{3}$$

$$h(z) + \varepsilon < g(z) \text{ for a.a. } z \in U \text{ (recall that } h \prec g). \tag{4}$$

We choose $\delta > 0$ so small that

$$\begin{aligned} & \left| \hat{\xi}(z) \left(|x|^{p(z)-2}x - |w|^{p(z)-2}w \right) \right| \\ & \leq \|\hat{\xi}\|_\infty \left| |x|^{p(z)-2}x - |w|^{p(z)-2}w \right| < \varepsilon \\ & \text{if } |x - w| < \delta, z \in \bar{U} \text{ (recall that } p \in C^{0,1}(\bar{\Omega}) \text{)}. \end{aligned} \tag{5}$$

Then we have for a.a. $z \in U$

$$\begin{aligned} & -\Delta_{p(z)}(u + \delta) + \hat{\xi}(z)|u + \delta|^{p(z)-2}(u + \delta) \\ & = -\Delta_{p(z)}u + \hat{\xi}(z)|u + \delta|^{p(z)-2}(u + \delta) \\ & = h(z) + \hat{\xi}(z) \left[|u + \delta|^{p(z)-2}(u + \delta) - |u|^{p(z)-2}u \right] \\ & \leq h(z) + \varepsilon \text{ (see (5))} \\ & < g(z) \text{ (see (4))} \\ & = -\Delta_{p(z)}v + \hat{\xi}(z)v^{p(z)-1}, \\ & \Rightarrow u(z) + \delta \leq v(z) \text{ for all } z \in \bar{U} \\ & \quad \text{(by the weak comparison principle, see (3)),} \\ & \Rightarrow u(z) < v(z) \text{ for all } z \in \bar{U}, \\ & \Rightarrow E = \emptyset. \end{aligned}$$

Also, by the anisotropic maximum principle of Zhang [26], we have

$$\begin{aligned} & \frac{\partial y}{\partial n} \Big|_{\partial\Omega} > 0, \\ & \Rightarrow \frac{\partial(v - u)}{\partial n} \Big|_{\partial\Omega} < 0. \end{aligned}$$

The proof is now complete. \square

For the second strong comparison principle, we use the following open cone in $C^1(\bar{\Omega})$

$$D_+ = \{u \in C^1(\bar{\Omega}) : u(z) > 0 \text{ for all } z \in \Omega, u \Big|_{\partial\Omega \cap U^{-1}(0)} < 0\}.$$

Proposition 2.5. *If $p \in C^{0,1}(\bar{\Omega})$, $1 < p(z)$ for all $z \in \bar{\Omega}$, $\hat{\xi}, h, g \in L^\infty(\Omega)$, $\hat{\xi}(z) \geq 0$ for a.a. $z \in \Omega$, $0 < \eta \leq g(z) - h(z)$ for a.a. $z \in \Omega$ and $u, v \in C^1(\bar{\Omega})$ satisfy $u \leq v$ and*

$$\begin{aligned} & -\Delta_{p(z)}u + \hat{\xi}(z)|u|^{p(z)-2}u = h(z) \text{ in } \Omega, \\ & -\Delta_{p(z)}v + \hat{\xi}(z)|v|^{p(z)-2}v = g(z) \text{ in } \Omega, \end{aligned}$$

then $v - u \in D_+$.

Proof. The reasoning is similar to that in the proof of the previous proposition.

Let $w = v - u \geq 0$, $w \in C^1(\bar{\Omega})$. As in the proof of Proposition 2.4, we have

$$-\operatorname{div}(A(z)Dw) = g(z) - h(z) - \hat{\xi}(z) \left[|v|^{p(z)-2}v - |u|^{p(z)-2}u \right] \text{ in } \Omega. \tag{6}$$

In this case we have $a_{ij} \in W^{1,\infty}(\Omega)$ for all $i, j = 1, \dots, N$.

Suppose that for some $z_0 \in \Omega$, we have $w(z_0) = 0$. Then $u(z_0) = v(z_0)$. The map $(z, x) \mapsto |x|^{p(z)-2}x$ is uniformly continuous on $\overline{\Omega} \times \mathbb{R}$. So, we can find $\delta > 0$ small enough such that

$$g(z) - h(z) - \hat{\xi}(z) \left| |v(z)|^{p(z)-2}v(z) - |u(z)|^{p(z)-2}u(z) \right| \geq \frac{\eta}{2} > 0$$

for a.a. $z \in B_\delta(z_0) = \{z \in \Omega : |z - z_0| < \delta\}$.

From (6) we have

$$-\operatorname{div}(A(z)Dw) \geq \frac{\eta}{2} > 0 \text{ for a.a. } z \in B_\delta(z_0).$$

Invoking Theorem 4 of Vázquez [25], we have

$$w(z) > 0 \text{ for all } z \in B_\delta(z_0),$$

a contradiction to the fact that $w(z_0) = 0$. Therefore

$$w(z) > 0 \text{ for all } z \in \Omega.$$

Let $E_0 = \{z \in \partial\Omega : w(z) = 0\}$. We can assume that $E_0 \neq \emptyset$. Otherwise we already have that $v(z) > u(z)$ for all $z \in \overline{\Omega}$ and we are done. By Zhang [26] we have

$$\begin{aligned} \frac{\partial w}{\partial n}(z_0) &< 0, \\ \Rightarrow w &= v - u \in D_+. \end{aligned}$$

The proof is now complete. \square

Now we introduce our hypotheses on the data of problem (P_λ) .

H_0 : $p, q \in C^{0,1}(\overline{\Omega})$ and $1 < q_- \leq q(z) \leq q_+ < p_- \leq p(z) \leq p_+$ for all $z \in \overline{\Omega}$, $\xi \in L^\infty(\Omega)$.

We mention that the global regularity results of Fan [7] that we use are actually valid for $p \in C^{0,\beta}(\overline{\Omega})$ with $0 < \beta < 1$ (see assumption p_H in [7, p. 398]). On the other hand, the Poincaré inequality (that is also used here) is valid for $p(\cdot)$ being Lipschitz continuous. For this reason we have assumed that $p, q \in C^{0,1}(\overline{\Omega})$.

H_1 : $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a function which for all $x \in \mathbb{R}$, is measurable in $z \in \Omega$, for a.a. $z \in \Omega$ we have $f(z, \cdot) \in C^1(\mathbb{R})$ and

- (i) $0 \leq f(z, x) \leq a(z)(1 + x^{r(z)-1})$ for a.a. $z \in \Omega$, all $x \in \mathbb{R}$, with $a \in L^\infty(\Omega)$, $r \in C(\overline{\Omega})$ and $p_+ < r(z) < p(z)^*$ for all $z \in \overline{\Omega}$;
- (ii) if $F(z, x) = \int_0^x f(z, s)ds$, then $\lim_{x \rightarrow +\infty} \frac{F(z, x)}{x^{p_+}} = +\infty$ uniformly for a.a. $z \in \Omega$;
- (iii) if $e(z, x) = f(z, x)x - p_+F(z, x)$, then there exist $M > 0$ and $\hat{C} > \|\xi\|_\infty \left(\frac{p_+}{p_-} - 1\right)$ such that

$$e'_x(z, x) \geq \hat{C}x^{p(z)-1} \text{ for a.a. } z \in \Omega, \text{ all } x \geq M;$$

- (iv) $\lim_{x \rightarrow 0^+} \frac{f(z, x)}{x^{p_- - 1}} = 0$ uniformly for a.a. $z \in \Omega$.

Remark 2.6. Since we are looking for positive solutions and all the above hypotheses concern the positive semi-axis $\mathbb{R}_+ = [0, +\infty)$, without any loss of generality, we may assume that

$$f(z, x) = 0 \text{ for a.a. } z \in \Omega, \text{ all } x \leq 0. \tag{7}$$

Remark 2.7. Hypothesis $H_1(ii)$ implies that for a.a. $z \in \Omega$, the function $F(z, \cdot)$ is p_+ -superlinear. This combined with hypothesis $H_1(iii)$, says that for a.a. $z \in \Omega$, $f(z, \cdot)$ is $(p_+ - 1)$ -superlinear. However, the superlinearity of $f(z, \cdot)$ is not expressed using the usual for such problems AR -condition (see Ambrosetti and Rabinowitz [2]). Instead, we employ the less restrictive condition $H_1(iii)$ that permits the consideration

of $(p_+ - 1)$ -superlinear functions with “slower” growth near $+\infty$, which fail to satisfy the *AR*-condition (see the examples below). Note that hypothesis $H_1(iii)$ is not global and implies that for a.a. $z \in \Omega$, $e(z, \cdot)$ is eventually nondecreasing. Then Lemma 2.4(iv) of Li and Yang [15], implies that there exists $\mu \in L^1(\Omega)$ such that

$$e(z, x) \leq e(z, y) + \mu(z) \text{ for a.a. } z \in \Omega, \text{ all } 0 \leq x \leq y \tag{8}$$

(a global quasi-monotonicity condition on $e(z, \cdot)$). Also it is equivalent to saying that there exists $\hat{M} > 0$ such that for a.a. $z \in \Omega$, the function

$$x \mapsto \frac{f(z, x)}{x^{p_+-1}}$$

is nondecreasing on $[\hat{M}, +\infty)$ (see Li and Yang [15]).

Example 2.8. Consider the following two functions

$$f_1(z, x) = x^{r(z)-1} \text{ for all } x \geq 0,$$

$$f_2(z, x) = \begin{cases} \hat{C}x^{r(z)-1}, & \text{if } x \in [0, 1] \\ \hat{C}(x^{p(z)-1} + x^{m(z)-1}), & \text{if } 1 < x \end{cases} \text{ (see (7)).}$$

with $r, m \in C(\bar{\Omega})$, $p_+ < r(z) < p(z)^*$, $m(z) \leq p(z)$ and $r(z) = p(z) + m(z) - 2$.

Note that $f_1(z, \cdot)$ satisfies the *AR*-condition, while $f_2(z, \cdot)$ need not (the *AR*-condition is not satisfied if $\{z \in \bar{\Omega} : m(z) = p_+\}$ has nonempty interior).

Finally, we mention that if X is a Banach space and $\varphi \in C^1(X, \mathbb{R})$, then K_φ denotes the critical set of φ , that is,

$$K_\varphi = \{u \in X : \varphi'(u) = 0\}.$$

Moreover, a set $S \subseteq W_0^{1,p(z)}(\Omega)$ is said to be “downward directed”, if given $u, v \in S$, we can find $w \in S$ such that $w \leq u$, $w \leq v$. In addition if $u, v \in W_0^{1,p(z)}(\Omega)$ with $u \leq v$, then we define

$$[u, v] = \{y \in W_0^{1,p(z)}(\Omega) : u(z) \leq y(z) \leq v(z) \text{ for a.a. } z \in \Omega\}$$

$$\text{int}_{C_0^1(\bar{\Omega})}[u, v] = \text{the interior in } C_0^1(\bar{\Omega}) \text{ of } [u, v] \cap C_0^1(\bar{\Omega}),$$

$$[u) = \{y \in W_0^{1,p(z)}(\Omega) : u(z) \leq y(z) \text{ for a.a. } z \in \Omega\}.$$

3. Positive solutions

We introduce the following two sets:

$$\mathcal{L} = \{\lambda > 0 : \text{problem } (P_\lambda) \text{ admits a positive solution}\},$$

$$S_\lambda = \text{set of positive solutions of problem } (P_\lambda).$$

Let $\vartheta > \|\xi\|_\infty$ (see hypothesis H_0), $\lambda > 0$ and consider the functional $\hat{\varphi}_\lambda : W_0^{1,p(z)}(\Omega) \rightarrow \mathbb{R}$ defined by

$$\hat{\varphi}_\lambda(u) = \int_\Omega \frac{1}{p(z)} |Du|^{p(z)} dz + \int_\Omega \frac{\xi(z)}{p(z)} |u|^{p(z)} dz + \int_\Omega \frac{\vartheta}{p(z)} (u^-)^{p(z)} dz$$

$$- \lambda \int_\Omega \frac{1}{q(z)} (u^+)^{q(z)} dz - \int_\Omega F(z, u^+) dz \text{ for all } u \in W_0^{1,p(z)}(\Omega).$$

Recall that $u^+ = \max\{u, 0\}$, $u^- = \max\{-u, 0\}$ and if $u \in W_0^{1,p(z)}(\Omega)$, then

$$u^+, u^- \in W_0^{1,p(z)}(\Omega), u = u^+ - u^-, |u| = u^+ + u^-.$$

We have that $\hat{\varphi}_\lambda(\cdot) \in C^1(W_0^{1,p(z)}(\Omega))$ (see Rădulescu and Repovš [23, p. 31]).

Proposition 3.1. *If hypotheses H_0, H_1 hold and $\lambda > 0$, then $\hat{\varphi}_\lambda(\cdot)$ satisfies the C-condition.*

Proof. We consider a sequence $\{u_n\}_{n \geq 1} \subseteq W_0^{1,p(z)}(\Omega)$ such that

$$|\hat{\varphi}_\lambda(u_n)| \leq M_1 \text{ for some } M_1 > 0, \text{ all } n \in \mathbb{N}, \tag{9}$$

$$(1 + \|u_n\|)\hat{\varphi}'_\lambda(u_n) \rightarrow 0 \text{ in } W^{-1,p'(z)}(\Omega) = W_0^{1,p(z)}(\Omega)^* \text{ as } n \rightarrow \infty. \tag{10}$$

From (10) we have

$$\begin{aligned} |\langle \hat{\varphi}'_\lambda(u_n), h \rangle| &\leq \frac{\varepsilon_n \|h\|}{1 + \|u_n\|} \text{ for all } h \in W_0^{1,p(z)}(\Omega), \text{ all } n \in \mathbb{N}, \text{ with } \varepsilon_n \rightarrow 0^+, \\ \Rightarrow \left| \langle A(u_n), h \rangle + \int_\Omega \xi(z) |u_n|^{p(z)-2} u_n h dz - \int_\Omega \vartheta (u_n^-)^{p(z)-1} h dz \right. \\ &\quad \left. - \lambda \int_\Omega (u_n^+)^{q(z)-1} h dz - \int_\Omega f(z, u_n^+) h dz \right| \leq \frac{\varepsilon_n \|h\|}{1 + \|u_n\|} \\ &\text{for all } h \in W_0^{1,p(z)}(\Omega), n \in \mathbb{N}. \end{aligned} \tag{11}$$

In (11) we choose $h = -u_n^- \in W_0^{1,p(z)}(\Omega)$. We have

$$\begin{aligned} \left| \hat{\rho}(Du_n^-) + \int_\Omega [\xi(z) + \vartheta] (u_n^-)^{p(z)} dz \right| &\leq \varepsilon_n \text{ for all } n \in \mathbb{N}, \\ \Rightarrow u_n^- &\rightarrow 0 \text{ in } W_0^{1,p(z)}(\Omega) \\ &\text{(recall that } \vartheta > \|\xi\|_\infty \text{ and see Proposition 2.2(d)).} \end{aligned} \tag{12}$$

In (11) we choose $h = u_n^+ \in W_0^{1,p(z)}(\Omega)$. Then

$$\left| \hat{\rho}(Du_n^+) + \int_\Omega \xi(z) (u_n^+)^{p(z)} dz - \lambda \int_\Omega (u_n^+)^{q(z)} dz - \int_\Omega f(z, u_n^+) u_n^+ dz \right| \leq \varepsilon_n \tag{13}$$

for all $n \in \mathbb{N}$.

On the other hand, from (9) and (12), we have

$$\begin{aligned} \left| \int_\Omega \frac{p_+}{p(z)} |Du_n^+|^{p(z)} dz + \int_\Omega \frac{p_+}{p(z)} \xi(z) |u_n^+|^{p(z)} dz - \lambda \int_\Omega \frac{p_+}{q(z)} (u_n^+)^{q(z)} dz \right. \\ \left. - \int_\Omega p_+ F(z, u_n^+) dz \right| \leq M_2 \\ \text{for some } M_2 > 0, \text{ all } n \in \mathbb{N}. \end{aligned} \tag{14}$$

From (13) and (14) it follows that

$$\begin{aligned} \int_\Omega \left[\frac{p_+}{p(z)} - 1 \right] |Du_n^+|^{p(z)} dz + \int_\Omega \left[\frac{p_+}{p(z)} - 1 \right] \xi(z) (u_n^+)^{p(z)} dz \\ - \lambda \int_\Omega \left[\frac{p_+}{q(z)} - 1 \right] (u_n^+)^{q(z)} dz + \int_\Omega e(z, u_n^+) dz \leq M_3 \\ \text{for some } M_3 > 0, \text{ all } n \in \mathbb{N}. \end{aligned} \tag{15}$$

Let $\beta_\lambda(z, x) = \lambda \left[1 - \frac{p_+}{q(z)} \right] x^{q(z)} + e(z, x) + \xi(z) \left[\frac{p_+}{p(z)} - 1 \right] x^{p(z)}$ for all $x \geq 0$.

Then from (15) we have

$$\int_\Omega \beta_\lambda(z, u_n^+) dz \leq M_3 \text{ for all } n \in \mathbb{N}. \tag{16}$$

Claim. The sequence $\{u_n^+\}_{n \geq 1} \subseteq W_0^{1,p(z)}(\Omega)$ is bounded.

We argue by contradiction. So, suppose that the claim is not true. Then passing to a subsequence if necessary, we may assume that

$$\|u_n^+\| \rightarrow \infty \text{ as } n \rightarrow \infty. \tag{17}$$

Let $y_n = \frac{u_n^+}{\|u_n^+\|}$, $n \in \mathbb{N}$. Then $\|y_n\| = 1$, $y_n \geq 0$ for all $n \in \mathbb{N}$. So, we may assume that

$$y_n \xrightarrow{w} y \text{ in } W_0^{1,p(z)}(\Omega) \text{ and } y_n \rightarrow y \text{ in } L^{p(z)}(\Omega), \ y \geq 0. \tag{18}$$

Let $\Omega_+ = \{z \in \Omega : y(z) > 0\}$ and $\Omega_0 = \{z \in \Omega : y(z) = 0\}$. Then $\Omega = \Omega_+ \cup \Omega_0$ (see (18)).

First we assume that $|\Omega_+|_N > 0$ (by $|\cdot|_N$ we denote the Lebesgue measure on \mathbb{R}^N). We have $u_n^+(z) \rightarrow +\infty$ for a.a. $z \in \Omega_+$ and so on account of hypothesis $H_1(ii)$ we have

$$\begin{aligned} & \frac{F(z, u_n^+(z))}{u_n^+(z)^{p_+}} \rightarrow +\infty \text{ for a.a. } z \in \Omega_+, \\ \Rightarrow & \frac{F(z, u_n^+(z))}{\|u_n^+\|^{p_+}} = \frac{F(z, u_n^+(z))}{u_n^+(z)^{p_+}} y_n(z)^{p_+} \rightarrow +\infty \text{ for a.a. } z \in \Omega_+, \\ \Rightarrow & \int_{\Omega_+} \frac{F(z, u_n^+)}{\|u_n^+\|^{p_+}} dz \rightarrow +\infty \text{ (by Fatou's lemma),} \\ \Rightarrow & \int_{\Omega} \frac{F(z, u_n^+)}{\|u_n^+\|^{p_+}} dz \rightarrow +\infty \text{ as } n \rightarrow +\infty. \end{aligned} \tag{19}$$

On account of (17), we may assume that $\|u_n^+\| \geq 1$ for all $n \in \mathbb{N}$. Then from (9) and (12), we have

$$\begin{aligned} & \lambda \int_{\Omega} \frac{1}{q(z)} \frac{(u_n^+)^{q(z)}}{\|u_n^+\|^{p_+}} dz + \int_{\Omega} \frac{F(z, u_n^+)}{\|u_n^+\|^{p_+}} dz \\ & \leq \varepsilon'_n + \frac{1}{p_-} \int_{\Omega} |Dy_n|^{p(z)} dz + \frac{\|\xi\|_{\infty}}{p_-} \int_{\Omega} y_n^{p(z)} dz \text{ with } \varepsilon'_n \rightarrow 0^+ \\ & \leq M_4 \text{ for some } M_4 > 0, \text{ all } n \in \mathbb{N} \text{ (see (18)).} \end{aligned} \tag{20}$$

Comparing (19) and (20), we have a contradiction.

So, we assume that $y \equiv 0$ (that is, $|\Omega|_N = |\Omega_0|_N$). We define

$$\hat{\varphi}_{\lambda}(t_n u_n) = \max\{\hat{\varphi}_{\lambda}(t u_n) : 0 \leq t \leq 1\}. \tag{21}$$

Let $v_n = \eta^{\frac{1}{p_-}} y_n$ for all $n \in \mathbb{N}$, with $\eta > 0$. Evidently we have

$$v_n \xrightarrow{w} 0 \text{ in } W_0^{1,p(z)}(\Omega) \text{ (see (18)).} \tag{22}$$

Hypothesis $H_1(i)$, (18) and the dominated convergence theorem imply that

$$\int_{\Omega} F(z, v_n) dz \rightarrow 0 \text{ as } n \rightarrow \infty. \tag{23}$$

Also, we have

$$\begin{aligned} & \int_{\Omega} \frac{1}{p(z)} \xi(z) v_n^{p(z)} dz \rightarrow 0, \int_{\Omega} \frac{1}{q(z)} v_n^{q(z)} dz \rightarrow 0 \\ & \text{(see (22) and Proposition 2.1).} \end{aligned} \tag{24}$$

Moreover, (17) implies that we can find $n_0 \in \mathbb{N}$ such that

$$\frac{\eta^{\frac{1}{p_-}}}{\|u_n^+\|} \in (0, 1] \text{ for all } n \geq n_0. \tag{25}$$

Then from (21) and (25), we have

$$\begin{aligned} \hat{\varphi}_\lambda(t_n u_n^+) &\geq \hat{\varphi}_\lambda(v_n) \\ &= \int_\Omega \frac{1}{p(z)} |Dv_n|^{p(z)} dz + \int_\Omega \frac{1}{p(z)} \xi(z) v_n^{p(z)} dz \\ &\quad - \lambda \int_\Omega \frac{1}{q(z)} v_n^{q(z)} dz - \int_\Omega F(z, v_n) dz \text{ for all } n \geq n_0, \\ &\geq \frac{1}{2p_+} \eta \text{ for all } n \geq n_1 \geq n_0 \\ &\quad \text{(see (23), (24) and use the Poincaré inequality).} \end{aligned}$$

Since $\eta > 0$ is arbitrary, we can infer that

$$\hat{\varphi}_\lambda(t_n u_n^+) \rightarrow +\infty \text{ as } n \rightarrow \infty. \tag{26}$$

We know that

$$\hat{\varphi}_\lambda(0) = 0 \text{ and } \hat{\varphi}_\lambda(u_n^+) \leq M_5 \text{ for some } M_5 > 0, \text{ all } n \in \mathbb{N}. \tag{27}$$

From (26) and (27) it follows that we can find $n_2 \in \mathbb{N}$ such that

$$t_n \in (0, 1) \text{ for all } n \geq n_2. \tag{28}$$

Then from (21) and (28) we infer that

$$\begin{aligned} t_n \frac{d}{dt} \hat{\varphi}_\lambda(tu_n^+) \Big|_{t=t_n} &= 0, \\ \Rightarrow \langle \hat{\varphi}'_\lambda(t_n u_n^+), t_n u_n^+ \rangle &= 0 \text{ for all } n \geq n_2 \\ &\text{(by the chain rule).} \end{aligned} \tag{29}$$

For all $n \geq n_2$ we have

$$\begin{aligned} &\hat{\varphi}_\lambda(t_n u_n^+) \\ &= \hat{\varphi}_\lambda(t_n u_n^+) - \frac{1}{p_+} \langle \hat{\varphi}'_\lambda(t_n u_n^+), t_n u_n^+ \rangle \text{ (see (29))} \\ &\leq \int_\Omega \left[\frac{1}{p(z)} - \frac{1}{p_+} \right] |D(t_n u_n^+)|^{p(z)} dz + \int_\Omega \left[\frac{1}{p(z)} - \frac{1}{p_+} \right] \xi(z) (t_n u_n^+)^{p(z)} dz \\ &\quad - \lambda \int_\Omega \left[\frac{1}{q(z)} - \frac{1}{p_+} \right] (t_n u_n^+)^{q(z)} dz + \frac{1}{p_+} \int_\Omega e(z, t_n u_n^+) dz \\ &\leq \int_\Omega \left[\frac{1}{p(z)} - \frac{1}{p_+} \right] |Du_n^+|^{p(z)} dz + \frac{1}{p_+} \int_\Omega \beta_\lambda(z, t_n u_n^+) dz. \end{aligned} \tag{30}$$

For the integrand $\beta_\lambda(z, x)$, we have for a.a. $z \in \Omega$, all $x \geq 0$

$$\begin{aligned} (\beta_\lambda)'_x(z, x) &= \lambda [q(z) - p_+] x^{q(z)-1} + e'_x(z, x) + \xi(z) [p_+ - p(z)] x^{p(z)-1} \\ &\geq \hat{C} x^{p(z)-1} - \lambda C_1 x^{q(z)-1} \text{ for some } C_1 > 0 \\ &\text{(see hypothesis } H_1(iii)). \end{aligned}$$

Since $q_+ < p_-$, we can find $M_6 \geq 1$ such that

$$\begin{aligned} &(\beta_\lambda)'_x(z, x) \geq 0 \text{ for a.a. } z \in \Omega, \text{ all } x \geq M_6, \\ \Rightarrow \beta_\lambda(z, \cdot) &\text{ is nondecreasing on } [M_6, \infty) \text{ for a.a. } z \in \Omega, \end{aligned}$$

$$\begin{aligned} \Rightarrow \beta_\lambda(z, x) &\leq \beta_\lambda(z, y) + \mu_\lambda(z) \\ &\text{for a.a. } z \in \Omega, \text{ all } 0 \leq x \leq y, \text{ with } \mu_\lambda \in L^1(\Omega). \end{aligned} \quad (31)$$

It follows from (30) and (31) that

$$\begin{aligned} &\hat{\varphi}_\lambda(t_n u_n^+) \\ &\leq \int_\Omega \left[\frac{1}{p(z)} - \frac{1}{p_+} \right] |Du_n^+|^{p(z)} dz + \frac{1}{p_+} \int_\Omega \beta_\lambda(z, u_n^+) dz + \frac{1}{p} \|\mu_\lambda\|_1 \\ &\quad \text{for all } n \geq n_2 \\ &= \hat{\varphi}_\lambda(u_n^+) - \frac{1}{p_+} \langle \hat{\varphi}'_\lambda(u_n^+), u_n^+ \rangle \\ &\leq \hat{\varphi}_\lambda(u_n^+) + \frac{\varepsilon_n}{p_+} \text{ for all } n \geq n_2 \text{ (see (13)),} \\ &\Rightarrow \hat{\varphi}_\lambda(u_n^+) \rightarrow +\infty \text{ as } n \rightarrow \infty \text{ (see (26)),} \\ &\quad \text{a contradiction (see (27)).} \end{aligned}$$

Therefore $\{u_n^+\} \subseteq W_0^{1,p(z)}(\Omega)$ is bounded and this proves the claim.

Then from (12) and the claim it follows that

$$\{u_n\}_{n \geq 1} \subseteq W_0^{1,p(z)}(\Omega) \text{ is bounded.}$$

We may assume that

$$u_n \xrightarrow{w} u \text{ in } W_0^{1,p(z)}(\Omega) \text{ and } u_n \rightarrow u \text{ in } L^{r(z)}(\Omega) \text{ as } n \rightarrow \infty. \quad (32)$$

In (11) we choose $h = u_n - u \in W_0^{1,p(z)}(\Omega)$, pass to the limit as $n \rightarrow \infty$ and use (32). Then

$$\begin{aligned} &\lim_{n \rightarrow \infty} \langle A(u_n), u_n - u \rangle = 0, \\ &\Rightarrow u_n \rightarrow u \text{ in } W_0^{1,p(z)}(\Omega) \text{ (see Proposition 2.3),} \\ &\Rightarrow \hat{\varphi}_\lambda(\cdot) \text{ satisfies the } C\text{-condition.} \end{aligned}$$

The proof is now complete. \square

Proposition 3.2. *If hypotheses H_0, H_1 hold, then $\mathcal{L} \neq \emptyset$ and we have $S_\lambda \subseteq \text{int } C_+$ for every $\lambda \in \mathcal{L}$.*

Proof. On account of hypotheses $H_1(i), (iv)$, we see that given $\varepsilon > 0$, we can find $C_2 = C_2(\varepsilon) > 0$ such that

$$F(z, x) \leq \frac{\varepsilon}{p_+} x^{p_+} + C_2 x^{r_+} \text{ for a.a. } z \in \Omega, \text{ all } x \geq 0. \quad (33)$$

For every $u \in W_0^{1,p(z)}(\Omega)$, we have

$$\begin{aligned} \hat{\varphi}_\lambda(u) &\geq \int_\Omega \frac{1}{p(z)} |Du|^{p(z)} dz + \int_\Omega \frac{1}{p(z)} \xi(z) |u|^{p(z)} dz \\ &\quad - \frac{\lambda}{q_+} \int_\Omega (u^+)^{q(z)} dz - \frac{\varepsilon}{p_+} \|u\|_{p_+}^{p_+} - C_3 \|u\|^{r_+} \\ &\quad \text{for some } C_3 > 0 \text{ (see (33) and recall that } \vartheta > \|\xi\|_\infty). \end{aligned} \quad (34)$$

For $\|u\|_{p(z)} \leq 1$ we have

$$\begin{aligned} & \int_{\Omega} \frac{1}{p(z)} \xi(z) |u|^{p(z)} dz \leq \frac{\|\xi\|_{\infty}}{p_-} \|u\|_{p(z)}^{p_-} \leq C_4 \|u\|^{p_-} \text{ for some } C_4 > 0, \\ \Rightarrow & \int_{\Omega} \frac{1}{p(z)} \xi(z) (u^+)^{q_+} dz \leq \lambda \|u\|^{q_+} + C_5 \|u\|^{r_+} \\ & \text{for some } C_5 = C_5(\lambda) > 0 \text{ (recall that } q_+ < p_- \leq p_+ < r_+ \text{)}. \end{aligned} \tag{35}$$

We return to (34) and use (35). Then for $u \in W_0^{1,p(z)}(\Omega)$ with $\max\{\|u\|, \|u\|_{p(z)}\} \leq 1$ we have

$$\begin{aligned} \hat{\varphi}_{\lambda}(u) & \geq \frac{1}{p_+} (1 - \varepsilon C_6) \|u\|^{p_+} - C_7 (\lambda \|u\|^{q_+} + \|u\|^{r_+}) \\ & \text{for some } C_6, C_7 > 0. \end{aligned}$$

We choose $\varepsilon \in (0, \frac{1}{C_6})$ and obtain

$$\begin{aligned} \hat{\varphi}_{\lambda}(u) & \geq C_8 \|u\|^{p_+} - C_7 (\lambda \|u\|^{q_+} + \|u\|^{r_+}) \text{ for some } C_8 > 0, \\ \Rightarrow \hat{\varphi}_{\lambda}(u) & \geq [C_8 - C_7 (\lambda \|u\|^{q_+ - p_+} + \|u\|^{r_+ - p_+})] \|u\|^{p_+}. \end{aligned} \tag{36}$$

Consider the function

$$k_{\lambda}(t) = \lambda t^{q_+ - p_+} + t^{r_+ - p_+} \text{ for all } t > 0.$$

Evidently, $k_{\lambda} \in C^1(0, +\infty)$ and since $q_+ < p_- \leq p_+ < r_+$ we have

$$k_{\lambda}(t) \rightarrow +\infty \text{ as } t \rightarrow 0^+ \text{ or } t \rightarrow +\infty.$$

So, we can find $t_0 > 0$ such that

$$\begin{aligned} k_{\lambda}(t_0) & = \min\{k_{\lambda}(t) : t > 0\}, \\ \Rightarrow k'_{\lambda}(t_0) & = 0, \\ \Rightarrow \lambda(p_+ - q_+)t_0^{q_+ - p_+ - 1} & = (r_+ - p_+)t_0^{r_+ - p_+ - 1}, \\ \Rightarrow t_0 & = \left[\frac{\lambda(p_+ - q_+)}{r_+ - p_+} \right]^{\frac{1}{r_+ - q_+}}. \end{aligned} \tag{37}$$

Then

$$\begin{aligned} k_{\lambda}(t_0) & = \lambda \frac{r_+ - p_+}{r_+ - q_+} \frac{(r_+ - p_+)^{\frac{p_+ - q_+}{r_+ - q_+}}}{(p_+ - q_+)^{\frac{p_+ - q_+}{r_+ - q_+}}} + \lambda \frac{r_+ - p_+}{r_+ - q_+} \frac{(p_+ - q_+)^{\frac{r_+ - p_+}{r_+ - q_+}}}{(r_+ - p_+)^{\frac{r_+ - p_+}{r_+ - q_+}}}, \\ \Rightarrow k_{\lambda}(t_0) & \rightarrow 0 \text{ as } \lambda \rightarrow 0^+. \end{aligned}$$

Let $C_0 > 0$ be such that $\|\cdot\|_{p(z)} \leq C_0 \|\cdot\|$. So, we can find $\lambda_0 > 0$ such that

$$0 < t_0 \leq \min \left\{ \frac{1}{C_0}, 1 \right\} \text{ and } k_{\lambda}(t_0) < \frac{C_8}{C_7} \text{ for all } \lambda \in (0, \lambda_0) \text{ (see (36), (37)).}$$

Then from (36) it follows that

$$\hat{\varphi}_{\lambda}(u) \geq \hat{m}_{\lambda} > 0 \text{ for all } \|u\| = t_0. \tag{38}$$

On account of superlinearity hypothesis $H_1(ii)$, for $u \in \text{int } C_+$, we have

$$\hat{\varphi}_{\lambda}(tu) \rightarrow -\infty \text{ as } t \rightarrow +\infty. \tag{39}$$

Then (38), (39) and Proposition 3.1, permit the use of the mountain pass theorem. Therefore for every $\lambda \in (0, \lambda_0)$ we can find $u_\lambda \in W_0^{1,p(z)}(\Omega)$ such that

$$u_\lambda \in K_{\hat{\varphi}_\lambda} \text{ and } 0 < \hat{m}_\lambda \leq \hat{\varphi}_\lambda(u_\lambda) \text{ (see (38)).} \quad (40)$$

From (40) we have $u_\lambda \neq 0$ (recall that $\hat{\varphi}_\lambda(0) = 0$) and

$$\langle \hat{\varphi}'_\lambda(u_\lambda), h \rangle = 0 \text{ for all } h \in W_0^{1,p(z)}(\Omega). \quad (41)$$

Choosing $h = -u_\lambda^- \in W_0^{1,p(z)}(\Omega)$, we obtain

$$\begin{aligned} & \int_\Omega \frac{1}{p(z)} |Du_\lambda^-|^{p(z)} dz + \int_\Omega \frac{\vartheta + \xi(z)}{p(z)} (u_\lambda^-)^{p(z)} dz = 0, \\ \Rightarrow & \frac{1}{p_+} [\hat{\rho}(Du_\lambda^-) + C_9 \rho(u_\lambda^-)] \leq 0 \text{ for some } C_9 > 0, \\ \Rightarrow & u_\lambda \geq 0, u_\lambda \neq 0. \end{aligned}$$

Then from (41) it follows that u_λ is a positive solution (P_λ). As before, the anisotropic regularity theory (see [7,8]) implies that

$$u_\lambda \in C_+ \setminus \{0\}.$$

We have

$$\begin{aligned} & -\Delta_{p(z)} u(z) + \xi(z) u(z)^{p(z)-1} \geq 0 \text{ for a.a. } z \in \Omega, \\ \Rightarrow & \Delta_{p(z)} u(z) \leq \|\xi\|_\infty u(z)^{p(z)-1} \text{ for a.a. } z \in \Omega, \\ \Rightarrow & u \in \text{int } C_+ \text{ (see Zhang [26]).} \end{aligned}$$

So, we have proved that $(0, \lambda_0) \subseteq \mathcal{L}$ and so $\mathcal{L} \neq \emptyset$. Moreover, we have $S_\lambda \subseteq \text{int } C_+$ for all $\lambda > 0$. \square

Next, we show that \mathcal{L} is an interval.

Proposition 3.3. *If hypotheses H_0, H_1 hold, $\lambda \in \mathcal{L}$ and $0 < \mu < \lambda$, then $u \in \mathcal{L}$ and given $u_\lambda \in S_\lambda$, we can find $u_\mu \in S_\mu$ such that $u_\mu \leq u_\lambda$.*

Proof. Since $\lambda \in \mathcal{L}$, we can find $u_\lambda \in S_\lambda \subseteq \text{int } C_+$. With $\vartheta > \|\xi\|_\infty$, we introduce the Carathéodory function $g_\mu(z, x)$ defined by

$$g_\mu(z, x) = \begin{cases} \mu(x^+)^{q(z)-1} + f(z, x^+) + \vartheta(x^+)^{p(z)-1}, & \text{if } x \leq u_\lambda(z) \\ \mu u_\lambda(z)^{q(z)-1} + f(z, u_\lambda(z)) + \vartheta u_\lambda(z)^{p(z)-1}, & \text{if } u_\lambda(z) < x. \end{cases} \quad (42)$$

We set $G_\mu(z, x) = \int_0^x g_\mu(z, s) ds$ and consider the C^1 -functional $\Psi_\mu : W_0^{1,p(z)}(\Omega) \rightarrow \mathbb{R}$ defined by

$$\Psi_\mu(u) = \int_\Omega \frac{1}{p(z)} |Du|^{p(z)} dz + \int_\Omega \frac{\vartheta + \xi(z)}{p(z)} |u|^{p(z)} dz - \int_\Omega G_\mu(z, u) dz$$

for all $u \in W_0^{1,p(z)}(\Omega)$.

Since $\vartheta > \|\xi\|_\infty$, from (42) it is clear that $\Psi_\mu(\cdot)$ is coercive. Also, using the fact that $W_0^{1,p(z)}(\Omega) \hookrightarrow L^{p(z)}(\Omega)$ compactly, we see that $\Psi_\mu(\cdot)$ is sequentially weakly lower semicontinuous. So, by the Weierstrass-Tonelli theorem, there exists $u_\mu \in W_0^{1,p(z)}(\Omega)$ such that

$$\Psi_\mu(u_\mu) = \inf \left\{ \Psi_\mu(u) : u \in W_0^{1,p(z)}(\Omega) \right\}. \quad (43)$$

Since $q_+ < p_-$, we see that

$$\begin{aligned} \Psi_\mu(u_\mu) < 0 &= \Psi_\mu(0), \\ \Rightarrow u_\mu &\neq 0. \end{aligned}$$

From (43) we have

$$\begin{aligned} \Psi'_\mu(u_\mu) &= 0, \\ \Rightarrow \langle A(u_\mu), h \rangle + \int_\Omega [\vartheta + \xi(z)] |u_\mu|^{p(z)-2} u_\mu h dz &= \int_\Omega g_\mu(z, u_\mu) h dz \\ \text{for all } h \in W_0^{1,p(z)}(\Omega). \end{aligned} \tag{44}$$

In (44) first we choose $h = -u_\mu^- \in W_0^{1,p(z)}(\Omega)$. We obtain

$$\begin{aligned} \hat{\rho}(Du_\mu^-) + C_{10}\rho(u_\mu^-) &\leq 0 \text{ for some } C_{10} > 0 \text{ (see (42)),} \\ \Rightarrow u_\mu &\geq 0, u_\mu \neq 0. \end{aligned}$$

Next, in (44) we choose $h = (u_\mu - u_\lambda)^+ \in W_0^{1,p(z)}(\Omega)$. We have

$$\begin{aligned} &\langle A(u_\mu), (u_\mu - u_\lambda)^+ \rangle + \int_\Omega [\vartheta + \xi(z)] u_\mu^{p(z)-1} (u_\mu - u_\lambda)^+ dz \\ &= \int_\Omega [\mu u_\lambda^{q(z)-1} + f(z, u_\lambda) + \vartheta u_\lambda^{p(z)-1}] (u_\mu - u_\lambda)^+ dz \text{ (see (42))} \\ &\leq \int_\Omega [\lambda u_\lambda u^{q(z)-1} + f(z, u_\lambda) + \vartheta u_\lambda^{p(z)-1}] (u_\mu - u_\lambda)^+ dz \text{ (since } \mu < \lambda) \\ &= \langle A(u_\lambda), (u_\mu - u_\lambda)^+ \rangle + \int_\Omega [\vartheta + \xi(z)] u_\lambda^{p(z)-1} (u_\mu - u_\lambda)^+ dz \text{ (since } u_\lambda \in S_\lambda). \end{aligned}$$

The monotonicity of $A(\cdot)$ (see Proposition 2.3) and the fact that $\vartheta > \|\xi\|_\infty$ imply that

$$\begin{aligned} u_\mu &\leq u_\lambda, \\ \Rightarrow u_\mu &\in [0, u_\lambda], u_\mu \neq 0, \\ \Rightarrow u_\mu &\in S_\mu \subseteq \text{int } C_+ \text{ (see (42) and (44)).} \end{aligned}$$

The proof is now complete. \square

So, according to Proposition 3.3, the solution multifunction $\lambda \mapsto S_\lambda$ has a kind of weak monotonicity property. We can improve this monotonicity property by adding one more condition on the perturbation $f(z, \cdot)$.

The new hypotheses on $f(z, x)$ are the following:

H_2 : $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a function which is measurable in $z \in \Omega$, for a.a. $z \in \Omega$ we have $f(z, \cdot) \in C^1(\mathbb{R})$, hypotheses $H_2(i) \rightarrow (iv)$ are the same as the corresponding hypotheses $H_1(i) \rightarrow (iv)$, and

(v) for every $\rho > 0$, there exists $\hat{\xi}_\rho > 0$ such that for a.a. $z \in \Omega$ the function

$$x \mapsto f(z, x) + \hat{\xi}_\rho x^{p(z)-1}$$

is nondecreasing on $[0, \rho]$.

Remark 3.4. This is a one-sided local Hölder condition on $f(z, \cdot)$. It is satisfied if for every $\rho > 0$, we can find $\hat{C}_\rho > 0$ such that $f'_x(z, x) \geq -\hat{C}_\rho x^{p(z)-1}$ for a.a. $z \in \Omega$ and all $0 \leq x \leq \rho$.

Proposition 3.5. *If hypotheses H_0, H_2 hold, $\lambda \in \mathcal{L}$, $u_\lambda \in S_\lambda \subseteq \text{int } C_+$ and $\mu \in (0, \lambda)$, then $\mu \in \mathcal{L}$ and we can find $u_\mu \in S_\mu \subseteq \text{int } C_+$ such that*

$$u_\lambda - u_\mu \in \text{int } C_+.$$

Proof. From Proposition 3.3 we know that $\mu \in \mathcal{L}$ and there exists $u_\mu \in S_\mu \subseteq \text{int } C_+$ such that

$$u_\lambda - u_\mu \in C_+ \setminus \{0\}. \tag{45}$$

Let $\rho = \|u_\lambda\|_\infty$ and let $\hat{\xi}_\rho > 0$ be as postulated by hypothesis $H_2(v)$. We can always assume that $\hat{\xi}_\rho > \|\xi\|_\infty$. Then we have

$$\begin{aligned} & -\Delta_{p(z)}u_\mu + [\xi(z) + \hat{\xi}_\rho]u_\mu^{p(z)-1} \\ &= \mu u_\mu^{q(z)-1} + f(z, u_\mu) + \hat{\xi}_\rho u_\mu^{p(z)-1} \\ &\leq \mu u_\lambda^{q(z)-1} + f(z, u_\lambda) + \hat{\xi}_\rho u_\lambda^{p(z)-1} \text{ (see (45) and hypothesis } H_2(v)) \\ &\leq \lambda u_\lambda^{q(z)-1} + f(z, u_\lambda) + \hat{\xi}_\rho u_\lambda^{p(z)-1} \text{ (since } \mu < \lambda) \\ &= -\Delta_{p(z)}u_\lambda + [\xi(z) + \hat{\xi}_\rho]u_\lambda^{p(z)-1}. \end{aligned} \tag{46}$$

Note that since $u_\lambda \in \text{int } C_+$ and $\mu < \lambda$, we have

$$0 \prec (\lambda - \mu)u_\lambda^{q(z)-1}. \tag{47}$$

Then from (46), (47) and Proposition 2.4, we can conclude that

$$u_\lambda - u_\mu \in \text{int } C_+.$$

The proof is now complete. \square

Next, we show that for every $\lambda \in \mathcal{L}$, the solution set S_λ has a smallest element (minimal positive solution). To this end, first we consider the following auxiliary problem

$$\begin{cases} -\Delta_{p(z)}u(z) + |\xi(z)||u(z)|^{p(z)-2}u(z) = \lambda|u(z)|^{q(z)-2}u(z) \text{ in } \Omega, \\ u|_{\partial\Omega} = 0, \lambda > 0, u > 0. \end{cases} \tag{48}$$

Proposition 3.6. *If hypotheses H_0 hold and $\lambda > 0$, then problem (48) admits a unique positive solution $\bar{u}_\lambda \in \text{int } C_+$.*

Proof. We consider the C^1 -functional $\gamma_\lambda : W_0^{1,p(z)}(\Omega) \rightarrow \mathbb{R}$ defined by

$$\gamma_\lambda(u) = \int_\Omega \frac{1}{p(z)}|Du|^{p(z)}dz + \int_\Omega \frac{|\xi(z)|}{p(z)}|u|^{p(z)}dz - \lambda \int_\Omega \frac{1}{q(z)}(u^+)^{q(z)}dz$$

for all $u \in W_0^{1,p(z)}(\Omega)$.

Evidently, $\gamma_\lambda(\cdot)$ is coercive (since $q_+ < p_-$) and sequentially weakly lower semicontinuous. So, we can find $\bar{u}_\lambda \in W_0^{1,p(z)}(\Omega)$ such that

$$\begin{aligned} \gamma_\lambda(\bar{u}_\lambda) &= \min \left\{ \gamma_\lambda(u) : u \in W_0^{1,p(z)}(\Omega) \right\} < 0 = \gamma_\lambda(0) \text{ (since } q_+ < p_-), \\ \Rightarrow \bar{u}_\lambda &\neq 0. \end{aligned}$$

We have

$$\begin{aligned} \gamma'_\lambda(\bar{u}_\lambda) &= 0, \\ \Rightarrow \langle A(\bar{u}_\lambda), h \rangle &+ \int_\Omega |\xi(z)||\bar{u}_\lambda|^{p(z)-2}\bar{u}_\lambda h dz = \lambda \int_\Omega (\bar{u}_\lambda^+)^{q(z)-1} h dz \end{aligned} \tag{49}$$

for all $h \in W_0^{1,p(z)}(\Omega)$.

In (49) we choose $h = -\bar{u}_\lambda^- \in W_0^{1,p(z)}(\Omega)$. Then

$$\begin{aligned} & \hat{\rho}(D\bar{u}_\lambda^-) + \int_\Omega |\xi(z)|(\bar{u}_\lambda^-)^{p(z)} dz = 0, \\ & \Rightarrow \bar{u}_\lambda \geq 0, \bar{u}_\lambda \neq 0, \\ & \Rightarrow \bar{u}_\lambda \text{ is a positive solution of (48) (see (49))}, \\ & \Rightarrow \bar{u}_\lambda \in C_+ \setminus \{0\} \text{ (anisotropic regularity theory)}. \end{aligned}$$

Therefore

$$\begin{aligned} & \Delta_{p(z)}\bar{u}_\lambda(z) \leq \|\xi\|_\infty \bar{u}_\lambda(z)^{p(z)-1} \text{ for a.a. } z \in \Omega, \\ & \Rightarrow \bar{u}_\lambda \in \text{int } C_+ \text{ (see Zhang [26])}. \end{aligned}$$

Next, we show that this positive solution of (48) is unique.

Suppose that \bar{v}_λ is another positive solution of (48). Again we have $\bar{v}_\lambda \in \text{int } C_+$. On account of Proposition 4.1.22 of Papageorgiou, Rădulescu and Repovš [20, p. 274], we have $\frac{\bar{u}_\lambda}{\bar{v}_\lambda}, \frac{\bar{v}_\lambda}{\bar{u}_\lambda} \in L^\infty(\Omega)$. So, we can apply Theorem 2.5 of Takač and Giacomoni [24] and get

$$\begin{aligned} 0 & \leq \int_\Omega \left[\frac{-\Delta_{p(z)}\bar{u}_\lambda}{\bar{u}_\lambda^{p_+ - 1}} + \frac{-\Delta_{p(z)}\bar{v}_\lambda}{\bar{v}_\lambda^{p_+ - 1}} \right] (\bar{u}_\lambda^{p_+} - \bar{v}_\lambda^{p_+}) dz \\ & = \int_\Omega \left[\lambda \left(\bar{u}_\lambda^{q(z)-p_+} - \bar{v}_\lambda^{q(z)-p_+} \right) - |\xi(z)| \left(\bar{u}_\lambda^{p(z)-p_+} - \bar{v}_\lambda^{p(z)-p_+} \right) \right] (\bar{u}_\lambda^{p_+} - \bar{v}_\lambda^{p_+}) dz, \\ & \Rightarrow \bar{u}_\lambda = \bar{v}_\lambda \text{ (since } q_+ < p_- \leq p(z)). \end{aligned}$$

Therefore the positive solution $\bar{u}_\lambda \in \text{int } C_+$ of problem (48) is unique. \square

This solution $\bar{u}_\lambda \in \text{int } C_+$ provides a lower bound for the solution set S_λ .

Proposition 3.7. *If hypotheses H_0, H_1 hold and $\lambda \in \mathcal{L}$, then $\bar{u}_\lambda \leq u$ for all $u \in S_\lambda$.*

Proof. Let $u \in S_\lambda \subseteq \text{int } C_+$ and consider the Carathéodory function $\beta_\lambda(z, x)$ defined by

$$\beta_\lambda(z, x) = \begin{cases} \lambda(x^+)^{q(z)-1}, & \text{if } x \leq u(z) \\ \lambda u(z)^{q(z)-1}, & \text{if } u(z) < x. \end{cases} \tag{50}$$

We set $B_\lambda(z, x) = \int_0^x \beta_\lambda(z, s) ds$ and consider the C^1 -functional $\tau_\lambda : W_0^{1,p(z)}(\Omega) \rightarrow \mathbb{R}$ defined by

$$\tau_\lambda(u) = \int_\Omega \frac{1}{p(z)} |Du|^{p(z)} dz + \int_\Omega \frac{|\xi(z)|}{p(z)} |u|^{p(z)} dz - \int_\Omega B_\lambda(z, u) dz$$

for all $u \in W_0^{1,p(z)}(\Omega)$.

From (50) we see that $\tau_\lambda(\cdot)$ is coercive. Also, it is sequentially weakly lower semicontinuous. So, we can find $\tilde{u}_\lambda \in W_0^{1,p(z)}(\Omega)$ such that

$$\begin{aligned} & \tau_\lambda(\tilde{u}_\lambda) = \min \left\{ \tau_\lambda(u) : u \in W_0^{1,p(z)}(\Omega) \right\} < 0 = \tau_\lambda(0) \text{ (since } q_+ < p_-), \\ & \Rightarrow \tilde{u}_\lambda \neq 0. \end{aligned}$$

We have

$$\begin{aligned} & \tau'_\lambda(\tilde{u}) = 0, \\ & \Rightarrow \langle A(\tilde{u}_\lambda), h \rangle + \int_\Omega |\xi(z)| |\tilde{u}_\lambda|^{p(z)-2} \tilde{u}_\lambda h dz = \int_\Omega \beta_\lambda(z, \tilde{u}_\lambda) h dz \end{aligned} \tag{51}$$

for all $h \in W_0^{1,p(z)}(\Omega)$.

In (51) we first choose $h = -\tilde{u}_\lambda^- \in W_0^{1,p(z)}(\Omega)$ and infer that

$$\tilde{u}_\lambda \geq 0, \quad \tilde{u}_\lambda \neq 0.$$

Next, in (51) we choose $h = (\tilde{u}_\lambda - u)^+ \in W_0^{1,p(z)}(\Omega)$. We have

$$\begin{aligned} & \langle A(\tilde{u}_\lambda), (\tilde{u}_\lambda - u)^+ \rangle + \int_\Omega |\xi(z)| \tilde{u}_\lambda^{p(z)-1} (\tilde{u}_\lambda - u)^+ dz \\ &= \int_\Omega \lambda u^{q(z)-1} (\tilde{u}_\lambda - u)^+ dz \quad (\text{see (50)}) \\ &\leq \int_\Omega [\lambda u^{q(z)-1} + f(z, u)] (\tilde{u}_\lambda - u)^+ dz \quad (\text{since } f \geq 0) \\ &\leq \langle A(u), (\tilde{u}_\lambda - u)^+ \rangle + \int_\Omega |\xi(z)| u^{p(z)-1} (\tilde{u}_\lambda - u)^+ dz \quad (\text{since } u \in S_\lambda), \\ &\Rightarrow \tilde{u}_\lambda \leq u. \end{aligned}$$

So, we have proved that

$$\tilde{u}_\lambda \in [0, u] \setminus \{0\}. \quad (52)$$

Then it follows from (51), (52) and (50) that

$$\begin{aligned} & \tilde{u}_\lambda \text{ is a positive solution of (48),} \\ & \Rightarrow \tilde{u}_\lambda = \bar{u}_\lambda \in \text{int } C_+ \quad (\text{see Proposition 3.6}), \\ & \Rightarrow \bar{u}_\lambda \leq u \text{ for all } u \in S_\lambda. \end{aligned}$$

The proof is now complete. \square

Remark 3.8. Reasoning as in the above proof, we show that $\lambda \mapsto \bar{u}_\lambda$ is increasing that is, if $0 < \mu < \lambda$, then $\bar{u}_\lambda - \bar{u}_\mu \in C_+ \setminus \{0\}$.

We know that S_λ is downward directed (see Filippakis and Papageorgiou [9] and Papageorgiou, Rădulescu and Repovš [18] and recall that $A(\cdot)$ is monotone, see Proposition 2.3).

Proposition 3.9. *If hypotheses H_0, H_1 hold and $\lambda \in \mathcal{L}$, then there exists $u_\lambda^* \in S_\lambda \subseteq \text{int } C_+$ such that*

$$u_\lambda^* \leq u \text{ for all } u \in S_\lambda$$

(minimal positive solution of (P_λ)).

Proof. By Lemma 3.10 of Hu and Papageorgiou [14, p. 178], we know that we can find $\{u_n\}_{n \geq 1} \subseteq S_\lambda \subseteq \text{int } C_+$ decreasing (recall that S_λ is downward directed) such that

$$\inf_{n \geq 1} u_n = \inf S_\lambda.$$

Since $\bar{u}_\lambda \leq u_n \leq u_1$ for all $n \in \mathbb{N}$ (see Proposition 3.7), from hypothesis $H_1(i)$ it follows that

$$\{u_n\}_{n \geq 1} \subseteq W_0^{1,p(z)}(\Omega) \text{ is bounded.}$$

So, we may assume that

$$u_n \xrightarrow{w} u_\lambda^* \text{ in } W_0^{1,p(z)}(\Omega) \text{ and } u_n \rightarrow u_\lambda^* \text{ in } L^{r(z)}(\Omega) \text{ as } n \rightarrow \infty. \quad (53)$$

We have

$$\begin{aligned}
 & \langle A(u_n), u_n - u_\lambda^* \rangle + \int_\Omega \xi(z) u_n^{p(z)-1} (u_n - u_\lambda^*) dz \\
 &= \lambda \int_\Omega u_n^{q(z)-1} (u_n - u_\lambda^*) dz + \int_\Omega f(z, u_n) (u_n - u_\lambda^*) dz, \\
 &\Rightarrow \lim_{n \rightarrow \infty} \langle A(u_n), u_n - u_\lambda^* \rangle = 0, \\
 &\Rightarrow u_n \rightarrow u_\lambda^* \text{ in } W_0^{1,p(z)}(\Omega) \text{ (see Proposition 2.3).}
 \end{aligned} \tag{54}$$

Note that

$$\bar{u}_\lambda \leq u_\lambda^* \text{ and so } u_\lambda^* \neq 0,$$

$$\begin{aligned}
 \langle A(u_\lambda^*), h \rangle + \int_\Omega \xi(z) (u_\lambda^*)^{p(z)-1} h dz &= \lambda \int_\Omega (u_\lambda^*)^{q(z)-1} h dz + \int_\Omega f(z, u_\lambda^*) h dz \\
 \text{for all } h \in W_0^{1,p(z)}(\Omega) &\text{ (see (54)).}
 \end{aligned}$$

It follows that

$$u_\lambda^* \in S_\lambda \subseteq \text{int } C_+ \text{ and } u_\lambda^* = \inf S_\lambda.$$

The proof is now complete. \square

We set $\lambda^* = \sup \mathcal{L}$.

Proposition 3.10. *If hypotheses H_0, H_2 hold, then $\lambda^* < \infty$.*

Proof. On account of hypotheses $H_0, H_2(iv)$ and since $q_+ < p_-$, we see that we can find $\hat{\lambda} > 0$ such that

$$\hat{\lambda} x^{q(z)-1} + f(z, x) - \xi(z) x^{p(z)-1} \geq 0 \text{ for a.a. } z \in \Omega, \text{ all } x \geq 0. \tag{55}$$

Let $\lambda > \hat{\lambda}$ and suppose that $\lambda \in \mathcal{L}$. Then we can find $u_\lambda \in S_\lambda \subseteq \text{int } C_+$. Let $\Omega_0 \subset\subset \Omega$ (that is, $\Omega_0 \subseteq \bar{\Omega}_0 \subseteq \Omega$) and assume that $\partial\Omega_0$ is a C^2 -manifold. We set $m_0 = \min_{\bar{\Omega}_0} u_\lambda > 0$ (recall that $u_\lambda \in \text{int } C_+$). Also, let $\hat{\xi}_\rho > \|\xi\|_\infty$. Let $m_0^\delta = m_0 + \delta$ for $\delta > 0$ small enough. We have

$$\begin{aligned}
 & -\Delta_{p(z)} m_0^\delta + [\xi(z) + \hat{\xi}_\rho] (m_0^\delta)^{p(z)-1} \\
 &\leq [\xi(z) + \hat{\xi}_\rho] m_0^{p(z)-1} + \chi(\delta) \text{ with } \chi(\delta) \rightarrow 0^+ \text{ as } \delta \rightarrow 0^+ \\
 &\leq \hat{\lambda} m_0^{q(z)-1} + f(z, m_0) + \hat{\xi}_\rho m_0^{p(z)-1} + \chi(\delta) \text{ (see (55))} \\
 &\leq \hat{\lambda} u_\lambda^{q(z)-1} + f(z, u_\lambda) + \hat{\xi}_\rho u_\lambda^{p(z)-1} + \chi(\delta) \text{ (see hypothesis } H_2(iv)) \\
 &\leq \lambda u_\lambda^{q(z)-1} + f(z, u_\lambda) + \hat{\xi}_\rho u_\lambda^{p(z)-1} - [\lambda - \hat{\lambda}] m_0^{p(z)-1} + \chi(\delta) \\
 &\leq -\Delta_{p(z)} u_\lambda + [\xi(z) + \hat{\xi}_\rho] u_\lambda^{p(z)-1} \text{ in } \Omega_0 \text{ for all } \delta \in (0, 1) \text{ small enough.}
 \end{aligned} \tag{56}$$

Note that for $\delta \in (0, 1)$ small enough, we have

$$(\lambda - \hat{\lambda}) m_0^{p(z)-1} - \chi(\delta) \geq \eta > 0.$$

Then from (56) and Proposition 2.5, we have

$$u_\lambda - m_0^\delta \in D_+ \text{ for all } \delta \in (0, 1) \text{ small enough,}$$

a contradiction. This means that $0 < \lambda^* \leq \hat{\lambda} < \infty$. \square

According to this proposition, we have

$$(0, \lambda^*) \subseteq \mathcal{L} \subseteq (0, \lambda^*]. \quad (57)$$

We will show that for all $\lambda \in (0, \lambda^*)$, we have at least two positive smooth solutions for problem (P_λ) . To do this we need to strengthen a little the hypotheses on $f(z, \cdot)$. The new conditions on $f(z, x)$ are the following:

H_3 : $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a function measurable in $z \in \Omega$, for a.a. $z \in \Omega$ $f(z, \cdot) \in C^1(\mathbb{R})$, hypotheses $H_3(i) \rightarrow (v)$ are the same as the corresponding hypotheses $H_2(i) \rightarrow (v) = H_1(i) \rightarrow (v)$ and

(vi) for every $m > 0$, there exists $\eta_m > 0$ such that

$$f(z, x) \geq \eta_m > 0 \text{ for a.a. } z \in \Omega, \text{ all } x \geq m.$$

Proposition 3.11. *If hypotheses H_0, H_3 hold and $\lambda \in (0, \lambda^*)$, then problem (P_λ) admits at least two positive solutions*

$$u_0, \hat{u} \in \text{int } C_+, \quad u_0 \neq \hat{u}.$$

Proof. Let $\eta \in (\lambda, \lambda^*)$. We have $\eta \in \mathcal{L}$ (see (57)) and so we can find $u_\eta \in S_\eta \subseteq \text{int } C_+$. Then according to Proposition 3.5, we can find $u_0 \in S_\lambda \subseteq \text{int } C_+$ such that

$$u_\eta - u_0 \in \text{int } C_+. \quad (58)$$

Recall that $\bar{u}_\lambda \leq u_0$ (see Proposition 3.7). Let $\rho = \|u_0\|_\infty$ and let $\hat{\xi}_\rho > 0$ be as postulated by hypothesis $H_3(v) = H_2(v)$. We can assume that $\hat{\xi}_\rho > \|\xi\|_\infty$. Then we have

$$\begin{aligned} & -\Delta_{p(z)} \bar{u}_\lambda + [\xi(z) + \hat{\xi}_\rho] \bar{u}_\lambda^{p(z)-1} \\ & \leq -\Delta_{p(z)} \bar{u}_\lambda + [|\xi(z)| + \hat{\xi}_\rho] \bar{u}_\lambda^{p(z)-1} \\ & = \lambda \bar{u}_\lambda^{q(z)-1} + \hat{\xi}_\rho \bar{u}_\lambda^{p(z)-1} \quad (\text{see Proposition 3.6}) \\ & \leq \lambda u_0^{q(z)-1} + f(z, \bar{u}_\lambda) + \hat{\xi}_\rho \bar{u}_\lambda^{p(z)-1} \quad (\text{recall that } f \geq 0) \\ & \leq \lambda u_0^{q(z)-1} + f(z, u_0) + \hat{\xi}_\rho u_0^{p(z)-1} \\ & \quad (\text{see Proposition 3.7 and hypothesis } H_3(v) = H_2(v)) \\ & = -\Delta_{p(z)} u_0 + [\xi(z) + \hat{\xi}_\rho] u_0^{p(z)-1} \quad (\text{since } u_0 \in S_\lambda). \end{aligned} \quad (59)$$

On account of hypothesis $H_3(vi)$ and since $\bar{u}_\lambda \in \text{int } C_+$, we see that

$$0 \prec f(\cdot, \bar{u}_\lambda(\cdot)).$$

Then (59) and Proposition 2.4 imply that

$$u_0 - \bar{u}_\lambda \in \text{int } C_+. \quad (60)$$

It follows from (58) and (60) that

$$u_0 \in \text{int}_{C^1(\bar{\Omega})} [\bar{u}_\lambda, u_\eta]. \quad (61)$$

As before, let $\vartheta > \|\xi\|_\infty$ and consider the Carathéodory function $k_\lambda(z, x)$ defined by

$$k_\lambda(z, x) = \begin{cases} \lambda \bar{u}_\lambda(z)^{q(z)-1} + f(z, \bar{u}_\lambda(z)) + \vartheta \bar{u}_\lambda(z)^{p(z)-1}, & \text{if } x < \bar{u}_\lambda(z) \\ \lambda x^{q(z)-1} + f(z, x) + \vartheta x^{p(z)-1}, & \text{if } \bar{u}_\lambda(z) \leq x \leq u_\eta(z) \\ \lambda u_\eta(z)^{q(z)-1} + f(z, u_\eta(z)) + \vartheta u_\eta(z)^{p(z)-1}, & \text{if } u_\eta(z) < x. \end{cases} \quad (62)$$

We set $K_\lambda(z, x) = \int_0^x k_\lambda(z, s)ds$ and consider the C^1 -functional $\tau_\lambda : W_0^{1,p(z)}(\Omega) \rightarrow \mathbb{R}$ defined by

$$\tau_\lambda(u) = \int_\Omega \frac{1}{p(z)} |Du|^{p(z)} dz + \int_\Omega \frac{\vartheta + \xi(z)}{p(z)} |u|^{p(z)} dz - \int_\Omega K_\lambda(z, u) dz$$

for all $u \in W_0^{1,p(z)}(\Omega)$.

From (62) and since $\vartheta > \|\xi\|_\infty$, we infer that $\tau_\lambda(\cdot)$ is coercive. Also it is sequentially weakly lower semicontinuous. So, we can find $\tilde{u}_0 \in W_0^{1,p(z)}(\Omega)$ such that

$$\begin{aligned} \tau_\lambda(\tilde{u}_0) &= \min\{\tau_\lambda(u) : u \in W_0^{1,p(z)}(\Omega)\}, \\ \Rightarrow \tau'_\lambda(\tilde{u}_0) &= 0, \\ \Rightarrow \langle \tau'_\lambda(\tilde{u}_0), h \rangle &= 0 \text{ for all } h \in W_0^{1,p(z)}(\Omega). \end{aligned}$$

Choosing $h = (\bar{u}_\lambda - \tilde{u}_0)^+$ and $h = (\tilde{u}_0 - u_\eta)^+$ and using (62), we show as before that

$$\tilde{u}_0 \in [\bar{u}_\lambda, u_\eta] \cap \text{int } C_+.$$

Therefore we may assume that $\tilde{u}_0 = u_0$ or otherwise we already have a second positive smooth solution and so we are done.

Next, we consider the Carathéodory function

$$\hat{k}_\lambda(z, x) = \begin{cases} \lambda \bar{u}_\lambda(z) q(z)^{-1} + f(z, \bar{u}_\lambda(z)) + \vartheta \bar{u}_\lambda(z) p(z)^{-1}, & \text{if } x \leq \bar{u}_\lambda(z) \\ \lambda x q(z)^{-1} + f(z, x) + \vartheta x p(z)^{-1}, & \text{if } \bar{u}_\lambda(z) < x. \end{cases} \tag{63}$$

We define $\hat{K}_\lambda(z, x) = \int_0^x \hat{k}_\lambda(z, s)ds$ and introduce the C^1 -functional $\hat{\tau}_\lambda : W_0^{1,p(z)}(\Omega) \rightarrow \mathbb{R}$ defined by

$$\hat{\tau}_\lambda(u) = \int_\Omega \frac{1}{p(z)} |Du|^{p(z)} dz + \int_\Omega \frac{\vartheta + \xi(z)}{p(z)} |u|^{p(z)} dz - \int_\Omega \hat{K}_\lambda(z, u) dz$$

for all $u \in W_0^{1,p(z)}(\Omega)$.

From (62) and (63) it is clear that

$$\tau_\lambda \Big|_{[\bar{u}_\lambda, u_\eta]} = \hat{\tau}_\lambda \Big|_{[\bar{u}_\lambda, u_\eta]}.$$

On account of (61), we have that

$$\begin{aligned} u_0 &\text{ is a local } C_0^1(\bar{\Omega})\text{-minimizer of } \hat{\tau}_\lambda, \\ \Rightarrow u_0 &\text{ is a local } W_0^{1,p(z)}(\Omega)\text{-minimizer of } \hat{\tau}_\lambda. \end{aligned} \tag{64}$$

(see Gasinski and Papageorgiou [11, Proposition 3.3]).

Using (63), we can easily see that

$$K_{\hat{\tau}_\lambda} \subseteq [\bar{u}_\lambda] \cap \text{int } C_+. \tag{65}$$

Then from (63) and (65) we can infer that we may assume that $K_{\hat{\tau}_\lambda}$ is finite or otherwise we already have an infinity of positive smooth solutions all distinct from u_0 and so, we are done. According to Theorem 5.7.6 of Papageorgiou, Rădulescu and Repovš [20, p. 449], we can find $\rho \in (0, 1)$ small such that

$$\hat{\tau}_\lambda(u_0) < \inf \{ \hat{\tau}_\lambda(u) : \|u - u_0\| = \rho \} = \hat{m}_\rho. \tag{66}$$

On account of hypothesis $H_3(ii) = H_1(ii)$, for $u \in \text{int } C_+$, we have

$$\hat{\tau}_\lambda(tu) \rightarrow -\infty \text{ as } t \rightarrow +\infty. \tag{67}$$

Finally, from (63) it follows that

$$\begin{aligned} \hat{\varphi}_\lambda \Big|_{[\bar{u}_\lambda]} &= \hat{\tau}_\lambda \Big|_{[\bar{u}_\lambda]} + \hat{\eta} \text{ with } \hat{\eta} \in \mathbb{R}, \\ \Rightarrow \hat{\tau}_\lambda(\cdot) &\text{ satisfies the } C\text{-condition (see Proposition 3.1)}. \end{aligned} \quad (68)$$

Then (66), (67), (68) permit the use of the mountain pass theorem. So, we can find $\hat{u} \in W_0^{1,p(z)}(\Omega)$ such that

$$\hat{u} \in K_{\hat{\tau}_\lambda} \subseteq [\bar{u}_\lambda] \cap \text{int } C_+ \text{ and } \hat{m}_\rho \leq \hat{\tau}_\lambda(\hat{u}). \quad (69)$$

From (69) and (63) we see that $\hat{u} \in S_\lambda \subseteq \text{int } C_+$, while from (69) and (66) we have that $\hat{u} \neq u_0$. \square

Finally, we show that the critical parameter value λ^* is admissible, that is, $\lambda^* \in \mathcal{L}$.

Proposition 3.12. *If hypotheses H_0, H_1 hold, then $\lambda^* \in \mathcal{L}$.*

Proof. Let $\{\lambda_n\}_{n \geq 1} \subseteq \mathcal{L}$ such that $\lambda_n \uparrow \lambda^*$ as $n \rightarrow \infty$. From the proof of Proposition 3.3, we know that we can find $u_n \in S_{\lambda_n} \subseteq \text{int } C_+$ such that

$$\hat{\varphi}_{\lambda_n}(u_n) < 0 \text{ for all } n \in \mathbb{N}.$$

Also, we have

$$\hat{\varphi}'_{\lambda_n}(u_n) = 0, \text{ for all } n \in \mathbb{N}.$$

Then as in the proof of Proposition 3.1, we show that

$$\{u_n\}_{n \geq 1} \subseteq W_0^{1,p(z)}(\Omega) \text{ is bounded.}$$

We may assume that

$$u_n \xrightarrow{w} u_* \text{ in } W_0^{1,p(z)}(\Omega) \text{ and } u_n \rightarrow u_* \text{ in } L^{r(z)}(\Omega) \text{ as } n \rightarrow \infty. \quad (70)$$

We have

$$\langle A(u_n), h \rangle + \int_{\Omega} \xi(z) u_n^{p(z)-1} h dz = \lambda_n \int_{\Omega} u_n^{q(z)-1} h dz + \int_{\Omega} f(z, u_n) h dz$$

for all $h \in W_0^{1,p(z)}(\Omega)$, all $n \in \mathbb{N}$.

Choosing $h = u_n - u_*$, passing to the limit as $n \rightarrow \infty$ and using (70) and Proposition 2.3, we obtain

$$u_n \rightarrow u_* \text{ in } W_0^{1,p(z)}(\Omega).$$

So, in the limit as $n \rightarrow \infty$, we have

$$\langle A(u_*), h \rangle + \int_{\Omega} \xi(z) u_*^{p(z)-1} h dz = \lambda^* \int_{\Omega} u_*^{q(z)-1} h dz + \int_{\Omega} f(z, u_*) h dz$$

for all $h \in W_0^{1,p(z)}(\Omega)$.

We have

$$\begin{aligned} \bar{u}_{\lambda_1} &\leq u_n \text{ for all } n \in \mathbb{N} \\ &\text{(see the Remark after Proposition 3.7),} \\ \Rightarrow \bar{u}_{\lambda_1} &\leq u_*, \\ \Rightarrow u_* &\in S_{\lambda^*} \subseteq \text{int } C_+ \text{ and so } \lambda^* \in \mathcal{L}. \end{aligned}$$

The proof is now complete. \square

According to this proposition, we have

$$\mathcal{L} = (0, \lambda^*].$$

Summarizing, we can state the following bifurcation-type result describing in a precise way the set of the positive solutions of problem (P_λ) as the parameter $\lambda > 0$ varies.

Theorem 3.13. If hypotheses H_0, H_3 hold, then there exists $\lambda^* > 0$ such that

(a) for all $\lambda \in (0, \lambda^*)$, problem (P_λ) has at least two positive solutions

$$u_0, \hat{u} \in \text{int } C_+, \quad u_0 \neq \hat{u};$$

(b) for $\lambda = \lambda^*$, problem (P_λ) has at least one positive solution

$$u_* \in \text{int } C_+;$$

(c) for $\lambda > \lambda^*$, problem (P_λ) has no positive solutions;

(d) for every $\lambda \in \mathcal{L} = (0, \lambda^*]$, problem (P_λ) has a smallest positive solution $u_\lambda^* \in \text{int } C_+$ and the map $\lambda \mapsto u_\lambda^*$ from $\mathcal{L} = (0, \lambda^*]$ into $C_+ \setminus \{0\}$ is increasing, that is,

$$0 < \mu \leq \lambda \in \mathcal{L} \Rightarrow u_\lambda^* - u_\mu^* \in C_+ \setminus \{0\}.$$

Acknowledgments

The authors wish to thank both anonymous reviewers for their remarks. This research was supported by the Slovenian Research Agency grants P1-0292, J1-8131, N1-0064, N1-0083, and N1-0114.

References

- [1] A. Ambrosetti, H. Brezis, G. Cerami, Combined effects of concave convex nonlinearities in some elliptic problems, *J. Funct. Anal.* 122 (1994) 519–543.
- [2] A. Ambrosetti, P. Rabinowitz, Dual variational methods in critical point theory and applications, *J. Funct. Anal.* 14 (1973) 349–381.
- [3] D. Arcoya, D. Ruiz, The Ambrosetti–Prodi problem for the p -Laplace operator, *Comm. Partial Differential Equations* 31 (2006) 849–865.
- [4] A. Bahrouni, V.D. Rădulescu, D.D. Repovš, A weighted anisotropic variant of the Caffarelli–Kohn–Nirenberg inequality and applications, *Nonlinearity* 31 (4) (2018) 1516–1534.
- [5] A. Bahrouni, V.D. Rădulescu, D.D. Repovš, Double phase transonic flow problems with variable growth: nonlinear patterns and stationary waves, *Nonlinearity* 32 (7) (2019) 2481–2495.
- [6] L. Diening, P. Harjulehto, P. Hästö, M. Ruzička, *Lebesgue and Sobolev Spaces with Variable Exponents*, in: *Lecture Notes in Mathematics*, vol. 2017, Springer, Heidelberg, 2011.
- [7] X. Fan, Global $C^{1,\alpha}$ regularity for variable exponent elliptic equations in divergence form, *J. Differential Equations* 235 (2007) 397–417.
- [8] X. Fan, D. Zhao, A class of De Giorgi type and Hölder continuity, *Nonlinear Anal.* 36 (1999) 295–318.
- [9] M. Filippakis, N.S. Papageorgiou, Multiple constant sign and nodal solutions for nonlinear elliptic equations with the p -Laplacian, *J. Differential Equations* 245 (2008) 1883–1929.
- [10] J. Garcia Azorero, J. Manfredi, I. Peral Alonso, Sobolev versus Hölder local minimizers and global multiplicity for some quasilinear elliptic equations, *Commun. Contemp. Math.* 2 (2000) 385–404.
- [11] L. Gasinski, N.S. Papageorgiou, Anisotropic nonlinear Neumann problems, *Calc. Var.* 42 (2011) 323–354.
- [12] M. Guedda, L. Véron, Quasilinear elliptic equations involving critical Sobolev exponents, *Nonlinear Anal.* 13 (1989) 879–902.
- [13] Z. Guo, Z. Zhang, $W^{1,p}$ Versus C^1 local minimizers and multiplicity results for quasilinear elliptic equations, *J. Math. Anal. Appl.* 286 (2003) 32–50.
- [14] S. Hu, N.S. Papageorgiou, *Handbook of Multivalued Analysis. Volume I: Theory*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1997.
- [15] G. Li, C. Yang, The existence of a nontrivial solution to a nonlinear elliptic boundary value problem of p -Laplacian type without Ambrosetti–Rabinowitz condition, *Nonlinear Anal.* 72 (2010) 4602–4613.

- [16] S. Marano, N.S. Papageorgiou, Positive solutions to a Dirichlet problem with p -Laplacian and concave-convex nonlinearity depending on a parameter, *Comm. Pure Appl. Anal.* 12 (2013) 815–829.
- [17] N.S. Papageorgiou, V.D. Rădulescu, Bifurcation of positive solutions for nonlinear nonhomogeneous Robin and Neumann problems with competing nonlinearities, *Discrete Contin. Dyn. Syst.* 35 (2016) 5008–5036.
- [18] N.S. Papageorgiou, V.D. Rădulescu, D.D. Repovš, Positive solutions for perturbations of the Robin eigenvalue problem plus an indefinite potential, *Discrete Contin. Dyn. Syst.* 37 (2017) 2589–2618.
- [19] N.S. Papageorgiou, V.D. Rădulescu, D.D. Repovš, Double-phase problems with reaction of arbitrary growth, *Z. Angew. Math. Phys.* 69 (4) (2018) 21, Art. 108.
- [20] N.S. Papageorgiou, V.D. Rădulescu, D.D. Repovš, *Nonlinear Analysis—Theory and Methods*, in: *Monographs in Mathematics*, Springer, Cham, 2019, Springer.
- [21] N.S. Papageorgiou, V.D. Rădulescu, D.D. Repovš, Double-phase problems and a discontinuity property of the spectrum, *Proc. Amer. Math. Soc.* 147 (7) (2019) 2899–2910.
- [22] N.S. Papageorgiou, V.D. Rădulescu, D.D. Repovš, Positive solutions for nonlinear parametric singular Dirichlet problems, *Bull. Math. Sci.* 9 (3) (2019) 21, 1950011.
- [23] V.D. Rădulescu, D.D. Repovš, *Partial Differential Equations with Variable Exponents: Variational Methods and Qualitative Analysis*, CRC Press, Boca Raton, FL, 2015.
- [24] P. Takač, J. Giacomoni, A $p(x)$ -Laplacian extension of the Diaz-Saa inequality and some applications, *Proc. Royal Soc. Edinburgh* (2020) <http://dx.doi.org/10.1017/prm.2018.91>.
- [25] J.-L. Vázquez, A strong maximum principle for some quasilinear elliptic equations, *Appl. Math. Optim.* 12 (1984) 191–202.
- [26] Q. Zhang, A strong maximum principle for differential equations with nonstandard $p(x)$ -growth conditions, *J. Math. Anal. Appl.* 312 (2005) 125–143.
- [27] Q. Zhang, V.D. Rădulescu, Double phase anisotropic variational problems and combined effects of reaction and absorption terms, *J. Math. Pures Appl.* 118 (2018) 159–203.